



# Lower Colorado River Multi-Species Conservation Program

*Balancing Resource Use and Conservation*

## Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL) Basic Conceptual Ecological Model for the Lower Colorado River



Photo courtesy of Robert Miller



February 2016

# Lower Colorado River Multi-Species Conservation Program Steering Committee Members

## **Federal Participant Group**

Bureau of Reclamation  
U.S. Fish and Wildlife Service  
National Park Service  
Bureau of Land Management  
Bureau of Indian Affairs  
Western Area Power Administration

## **Arizona Participant Group**

Arizona Department of Water Resources  
Arizona Electric Power Cooperative, Inc.  
Arizona Game and Fish Department  
Arizona Power Authority  
Central Arizona Water Conservation District  
Cibola Valley Irrigation and Drainage District  
City of Bullhead City  
City of Lake Havasu City  
City of Mesa  
City of Somerton  
City of Yuma  
Electrical District No. 3, Pinal County, Arizona  
Golden Shores Water Conservation District  
Mohave County Water Authority  
Mohave Valley Irrigation and Drainage District  
Mohave Water Conservation District  
North Gila Valley Irrigation and Drainage District  
Town of Fredonia  
Town of Thatcher  
Town of Wickenburg  
Salt River Project Agricultural Improvement and Power District  
Unit "B" Irrigation and Drainage District  
Wellton-Mohawk Irrigation and Drainage District  
Yuma County Water Users' Association  
Yuma Irrigation District  
Yuma Mesa Irrigation and Drainage District

## **Other Interested Parties Participant Group**

QuadState Local Governments Authority  
Desert Wildlife Unlimited

## **California Participant Group**

California Department of Fish and Wildlife  
City of Needles  
Coachella Valley Water District  
Colorado River Board of California  
Bard Water District  
Imperial Irrigation District  
Los Angeles Department of Water and Power  
Palo Verde Irrigation District  
San Diego County Water Authority  
Southern California Edison Company  
Southern California Public Power Authority  
The Metropolitan Water District of Southern California

## **Nevada Participant Group**

Colorado River Commission of Nevada  
Nevada Department of Wildlife  
Southern Nevada Water Authority  
Colorado River Commission Power Users  
Basic Water Company

## **Native American Participant Group**

Hualapai Tribe  
Colorado River Indian Tribes  
Chemehuevi Indian Tribe

## **Conservation Participant Group**

Ducks Unlimited  
Lower Colorado River RC&D Area, Inc.  
The Nature Conservancy



# Lower Colorado River Multi-Species Conservation Program

## Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL) Basic Conceptual Ecological Model for the Lower Colorado River

*Prepared by:*

Robert A. Miller, Christopher J.W. McClure, Ph.D., and  
Robert Unnasch, Ph.D.

Sound Science, LLC

Lower Colorado River  
Multi-Species Conservation Program  
Bureau of Reclamation  
Lower Colorado Region  
Boulder City, Nevada  
<http://www.lcrmscp.gov>

February 2016

Miller, R.A., C.J.W. McClure, and R. Unnasch. 2016. Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL) Basic Conceptual Ecological Model for the Lower Colorado River. Submitted to the Bureau of Reclamation, Boulder City, Nevada, by Sound Science, LLC, Boise, Idaho.

# ACRONYMS AND ABBREVIATIONS

CEM	conceptual ecological model
LCR	lower Colorado River
LCR MSCP	Lower Colorado River Multi-Species Conservation Program
Reclamation	Bureau of Reclamation
USFWS	U.S. Fish and Wildlife Service
VEFL	vermillion flycatcher

## Symbols

$\geq$	greater than or equal to
$<$	less than
$\pm$	plus or minus

## Definitions

For the purposes of this document, vegetation layers are defined as follows:

**Canopy** – The canopy is the uppermost strata within a plant community. The canopy is exposed to the sun and captures the majority of its radiant energy.

**Understory** – The understory comprises plant life growing beneath the canopy without penetrating it to any extent. The understory exists in the shade of the canopy and usually has lower light and higher humidity levels. The understory includes subcanopy trees and the shrub and herbaceous layers.

**Shrub layer** – The shrub layer is comprised of woody plants between 0.5 and 2.0 meters in height.

**Herbaceous layer** – The herbaceous layer is most commonly defined as the forest stratum composed of all vascular species that are 0.5 meter or less in height.

# CONTENTS

	Page
<b>Foreword</b> .....	v
<b>Executive Summary</b> .....	<b>ES-1</b>
Conceptual Ecological Models .....	ES-1
Conceptual Ecological Model Structure .....	ES-2
Results.....	ES-4
<b>Chapter 1 – Introduction</b> .....	<b>1</b>
Vermilion Flycatcher Reproductive Ecology .....	1
Conceptual Ecological Model Purposes .....	2
Conceptual Ecological Model Structure for the VEFL.....	3
<b>Chapter 2 – VEFL Life Stage Model</b> .....	<b>7</b>
Introduction to the VEFL Life Cycle.....	7
VEFL Life Stage 1 – Nest.....	8
VEFL Life Stage 2 – Juvenile.....	8
VEFL Life Stage 3 – Breeding Adult .....	8
Life Stage Model Summary .....	9
<b>Chapter 3 – Critical Biological Activities and Processes</b> .....	<b>11</b>
Disease .....	12
Eating/Foraging.....	12
Molt.....	12
Nest Attendance .....	13
Nest Site Selection .....	13
Predation and Brood Parasitism.....	13
Temperature Regulation.....	14
<b>Chapter 4 – Habitat Elements</b> .....	<b>15</b>
Anthropogenic Disturbance .....	16
Brood Size.....	17
Canopy Closure.....	17
Community Type .....	17
Food Availability .....	17
Genetic Diversity and Infectious Agents .....	18
Local Hydrology .....	18
Parental Feeding Behavior.....	19
Parental Nest Attendance .....	19
Patch Size.....	19
Predator and Cowbird Density.....	19
Soil Salinity.....	20
Tree Size .....	20

	Page
Understory Density .....	20
Water Availability.....	20
<b>Chapter 5 – Controlling Factors.....</b>	<b>21</b>
Fire Management .....	22
Grazing.....	22
Irrigation .....	22
Mechanical Thinning .....	22
Pesticide/Herbicide Application .....	23
Planting Regime.....	23
Recreational Activities.....	24
<b>Chapter 6 – Conceptual Ecological Model by Life Stage.....</b>	<b>25</b>
VEFL Life Stage 1 – Nest.....	28
VEFL Life Stage 2 – Juvenile.....	35
VEFL Life Stage 3 – Breeding Adult .....	41
<b>Chapter 7 – Causal Relationships Across All Life Stages.....</b>	<b>47</b>
Anthropogenic Disturbance .....	48
Canopy closure.....	48
Community Type .....	49
Food Availability .....	49
Local Hydrology .....	50
Parental Feeding Behavior.....	50
Parental Nest Attendance.....	50
Patch Size.....	51
Predator and Cowbird Density.....	52
Soil Salinity.....	52
Tree Size .....	52
Understory Density .....	53
Water Availability.....	54
<b>Chapter 8 – Discussion and Conclusions .....</b>	<b>55</b>
Most Influential Activities and Processes Across All Life Stages .....	55
Potentially Pivotal Alterations to Habitat Elements .....	56
Gaps in Understanding.....	57
Literature Cited .....	61
Acknowledgments.....	69

## Tables

Table		Page
ES-1	Outcomes of each of the three life stages of VEFL .....	3
1	VEFL life stages and outcomes in the LCR ecosystem .....	9
2	Distribution of VEFL critical biological activities and processes among life stages .....	11
3	Distribution of VEFL habitat elements and the critical biological activities and processes that they directly affect across all life stages .....	15
4	Habitat elements directly affected by controlling factors .....	21
5	Magnitude of influence of controlling factors on habitat elements ....	47

## Figures

Figure		Page
1	Proposed VEFL life history model. ....	10
2	Diagram conventions for LCR MSCP conceptual ecological models. ....	27
3	VEFL life stage 1 – nest, basic CEM diagram. Only elements with connections within this life stage are presented.....	31
4	VEFL life stage 1 – nest, high- and medium-magnitude relationships. Only elements with high- or medium-magnitude connections within this life stage are presented.....	33
5	VEFL life stage 2 – Juvenile, basic CEM diagram. Only elements with connections within this life stage are presented.....	37
6	VEFL life stage 2 – juvenile, high- and medium-magnitude relationships. Only elements with high- or medium-magnitude connections within this life stage are presented.....	39
7	VEFL life stage 3 – Breeding Adult, basic CEM diagram. Only elements with connections within this life stage are presented. ....	43
8	VEFL life stage 3 – breeding adult, high- and medium-magnitude relationships. Only elements with high- or medium-magnitude connections within this life stage are presented.....	45
9	Most influential biological activities and processes affecting each life stage of VEFL. ....	56
10	Habitat elements that directly affect the most influential biological activities and processes across all life stages of VEFL.....	58

## **Attachments**

### Attachment

- 1 Species Conceptual Ecological Model Methodology for the Lower Colorado River Multi-Species Conservation Program
- 2 Vermilion Flycatcher Habitat Data

# Foreword

The Lower Colorado River Multi-Species Conservation Program (LCR MSCP) Habitat Conservation Plan requires the creation, and long-term stewardship, of habitat for 20 covered species. This is both an exciting and daunting challenge – exciting, in that success would mean a major conservation achievement in the lower Colorado River landscape, and daunting, in that we need to simultaneously manage our lands for the benefit of 20 species in a mosaic of land cover types. To do so, we need to develop a common understanding of the habitat requirements of each species and the stewardship required to meet those needs.

To provide a framework to capture and share the information that forms the foundation of this understanding, conceptual ecological models (CEMs) for each covered species have been created under the LCR MSCP’s Adaptive Management Program. The LCR MSCP’s conceptual ecological models are descriptions of the functional relationships among essential components of a species’ life history, including its habitat, threats, and drivers. They tell the story of “what’s important to the animal” and how our stewardship and restoration actions can change those processes or attributes for the betterment of their habitat. As such, CEMs can provide:

- A synthesis of the current understanding of how a species’ habitat works. This synthesis can be based on the published literature, technical reports, or professional experience.
- Help in understanding and diagnosing underlying issues and identifying land management opportunities.
- A basis for isolating cause and effect and simplifying complex systems. These models also document the interaction among system drivers.
- A common (shared) framework or “mental picture” from which to develop management alternatives.
- A tool for making qualitative predictions of ecosystem responses to stewardship actions.
- A way to flag potential thresholds from which system responses may accelerate or follow potentially unexpected or divergent paths.
- A means by which to outline further restoration, research, and development and to assess different restoration scenarios.

- A means of identifying appropriate monitoring indicators and metrics.
- A basis for implementing adaptive management strategies.

Most natural resource managers rely heavily upon CEMs to guide their work, but few explicitly formulate and express the models so they can be shared, assessed, and improved. When this is done, these models provide broad utility for ecosystem restoration and adaptive management.

Model building consists of determining system parts, identifying the relationships that link these parts, specifying the mechanisms by which the parts interact, identifying missing information, and exploring the model's behavior (Heemskerk et al. 2003<sup>1</sup>). The model building process can be as informative as the model itself, as it reveals what is known and what is unknown about the connections and causalities in the systems under management.

It is important to note that CEMs are not meant to be used as prescriptive management tools but rather to give managers the information needed to help inform decisions. These models are conceptual and qualitative. They are not intended to provide precise, quantitative predictions. Rather, they allow us to virtually “tweak the system” free of the constraints of time and cost to develop a prediction of how a system might respond over time to a variety of management options; for a single species, a documented model is a valuable tool, but for 20 species, they are imperative. The successful management of multiple species in a world of competing interests (species versus species), potentially conflicting needs, goals, and objectives, long response times, and limited resources, these models can help land managers experiment from the safety of the desktop. Because quantitative data can be informative, habitat parameters that have been quantified in the literature are presented (in attachment 2) in this document for reference purposes.

These models are intended to be “living” documents that should be updated and improved over time. The model presented here should not be viewed as a definitive monograph of a species' life history but rather as a framework for capturing the knowledge and experience of the LCR MSCP's scientists and land stewards. While ideally the most helpful land management tool would be a definitive list of do's and don'ts, with exact specifications regarding habitat requirements that would allow us to engineer exactly what the species we care about need to survive and thrive, this is clearly not possible. The fact is, that despite years of active management, observation, and academic research on many of the LCR MSCP species of concern, there may not be enough data to support developing such detailed, prescriptive land management.

---

<sup>1</sup> Heemskerk, M., K. Wilson, and M. Pavao-Zuckerman. 2003. Conceptual models as tools for communication across disciplines. *Conservation Ecology* 7(3):8:  
<http://www.consecol.org/vol7/iss3/art8/>

The CEMs for species covered under the LCR MSCP are based on, and expand upon, methods developed by the Sacramento-San Joaquin Delta Ecosystem Restoration Program (ERP): [https://www.dfg.ca.gov/ERP/conceptual\\_models.asp](https://www.dfg.ca.gov/ERP/conceptual_models.asp). The ERP is jointly implemented by the California Department of Fish and Wildlife, U.S. Fish and Wildlife Service, and National Marine Fisheries Service. The Bureau of Reclamation (Reclamation) participates in this program. (See attachment 1 for an introduction to the CEM process.)

Many of the LCR MSCP covered species are migratory. These models only address the species' life history as it relates to the lower Colorado River and specifically those areas that are potentially influenced by LCR MSCP land management. The models DO NOT take into account ecological factors that influence the species at their other migratory locations.

Finally, in determining the spatial extent of the literature used in these models, the goals and objectives of the LCR MSCP were taken into consideration. For species whose range is limited to the Southwest, the models are based on literature from throughout the species' range. In contrast, for those species whose breeding range is continental (e.g., yellow-billed cuckoo) or west-wide, the models primarily utilize studies from the Southwest.

### **How to Use the Models**

There are three important elements to each CEM:

- (1) The narrative description of the species' various life stages, critical biological activities and processes, and associated habitat elements.
- (2) The figures that provide a visual snapshot of all the critical factors and causal links for a given life stage.
- (3) The associated workbooks. Each CEM has a workbook that includes a worksheet for each life stage.

This narrative document is a basic guide, meant to summarize information on the species' most basic habitat needs, the figures are a graphic representation of how these needs are connected, and the accompanying workbook is a tool for land managers to see how on-the-ground changes might potentially change outcomes for the species in question. Reading, evaluating, and using these CEMs requires that the reader understand all three elements; no single element provides all the pertinent information in the model. While it seems convenient to simply read the narrative, we strongly recommend the reader have the figures and workbook open and refer to them while reviewing this document.

It is also tempting to see these products, once delivered, as “final.” However, it is more accurate to view them as “living” documents, serving as the foundation for future work. Reclamation will update these products as new information is available, helping to inform land managers as they address the on-the-ground challenges inherent in natural resource management.

The knowledge gaps identified by these models are meant to serve only as an example of the work that could be done to further complete our understanding of the life history of the LCR MSCP covered species. However, this list can in no way be considered an exhaustive list of research needs. Additionally, while identifying knowledge gaps was an objective of this effort, evaluating the feasibility of addressing those gaps was not. Finally, while these models were developed for the LCR MSCP, the identified research needs and knowledge gaps reflect a current lack of understanding within the wider scientific community. As such, they may not reflect the current or future goals of the LCR MSCP. They are for the purpose of informing LCR MSCP decisionmaking but are in no way meant as a call for Reclamation to undertake research to fill the identified knowledge gaps.

*John Swett, Program Manager, LCR MSCP  
Bureau of Reclamation  
September 2015*

# Executive Summary

This document presents a conceptual ecological model (CEM) for the vermilion flycatcher (*Pyrocephalus rubinus*) (VEFL). The purpose of this model is to help the Bureau of Reclamation (Reclamation), Lower Colorado River Multi-Species Conservation Program (LCR MSCP), identify areas of scientific uncertainty concerning VEFL ecology, the effects of specific stressors, the effects of specific management actions aimed at species habitat restoration, and the methods used to measure VEFL habitat and population conditions. (Note: Attachment 1 provides an introduction to the CEM process. We recommend that those unfamiliar with this process read the attachment before continuing with this document.)

The identified research questions and gaps in scientific knowledge that are the result of this modeling effort serve as examples of topics the larger scientific community could explore to improve the overall understanding of the ecology of this species. These questions may or may not be relevant to the goals of the LCR MSCP. As such, they are not to be considered guidance for Reclamation or the LCR MSCP, nor are these knowledge gaps expected to be addressed under the program.

## CONCEPTUAL ECOLOGICAL MODELS

CEMs integrate and organize existing knowledge concerning: (1) what is known about an ecological resource, with what certainty, and the sources of this information, (2) critical areas of uncertain or conflicting science that demand resolution to better guide management planning and action, (3) crucial attributes to use while monitoring system conditions and predicting the effects of experiments, management actions, and other potential agents of change, and (4) how we expect the characteristics of the resource to change as a result of altering its shaping/controlling factors, including those resulting from management actions.

The CEM applied to the VEFL expands on the methodology developed for the Sacramento-San Joaquin River Delta Regional Ecosystem Restoration Implementation Plan (DiGennaro et al. 2012). The model distinguishes the major life stages or events through which the individuals of a species must pass to complete a full life cycle. It then identifies the factors that shape the likelihood that individuals in each life stage will survive to the next stage in the study area and thereby shapes the abundance, distribution, and persistence of the species in that area.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

Specifically, the VEFL conceptual ecological model has five core components:

- **Life stages** – These consist of the major growth stages and critical events through which the individual VEFL must pass in order to complete a full reproductive cycle.
- **Life-stage outcomes** – These consist of the biologically crucial outcomes of each life stage, including the number of individuals recruited to the next life stage or age class within a single life stage (recruitment rate), or the number of offspring produced (fertility rate).
- **Critical biological activities and processes** – These consist of activities in which the species engages and the biological processes that take place during each life stage that significantly beneficially or detrimentally shape the life-stage outcome rates for that life stage.
- **Habitat elements** – These consist of the specific habitat conditions, the abundance, spatial and temporal distributions, and other qualities that significantly beneficially or detrimentally affect the rates of the critical biological activities and processes for each life stage.
- **Controlling factors** – These consist of environmental conditions and dynamics – including human actions – that determine the abundance, spatial and temporal distributions, and other qualities of the habitat elements for each life stage. Controlling factors are also called “drivers.”

The CEM identifies the causal relationships among these components for each life stage. A causal relationship exists when a change in one condition or property of a system results in a change in some other condition or property. A change in the first condition is said to cause a change in the second condition. The CEM method applied here assesses four variables for each causal relationship: (1) the character and direction of the effect, (2) the magnitude of the effect, (3) the predictability (consistency) of the effect, and (4) the certainty of a present scientific understanding of the effect. CEM diagrams and a linked spreadsheet tool document all information on the model components and their causal relationships.

## **CONCEPTUAL ECOLOGICAL MODEL STRUCTURE**

The VEFL conceptual ecological model addresses the VEFL throughout its breeding range. The model thus addresses the landscape as a whole rather than any single reach or managed area. The model does not specifically address the biology of migratory VEFL during winter.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

The most widely used sources of the information for the VEFL conceptual ecological model are Rosenberg et al. (1991), Myers (2008), Ellison et al. (2009), Brand et al. (2010), and Raulston (2013). These publications summarize and cite large bodies of earlier studies. Where appropriate and accessible, those earlier studies are directly cited. The model also integrates numerous additional sources, particularly reports and articles completed since these publications, information on current research projects, and the expert knowledge of LCR MSCP biologists. Our purpose is not to provide an updated literature review, but to integrate the available information and knowledge into a CEM so it can be used for adaptive management.

The VEFL conceptual ecological model distinguishes and assesses three life stages and their associated outcomes as follows (table ES-1):

Table ES-1.—Outcomes of each of the three life stages of VEFL

<b>Life stage</b>	<b>Life-stage outcome(s)</b>
1. Nest	<ul style="list-style-type: none"><li>• Survival</li></ul>
2. Juvenile	<ul style="list-style-type: none"><li>• Survival</li></ul>
3. Breeding adult	<ul style="list-style-type: none"><li>• Survival</li><li>• Reproduction</li></ul>

The model distinguishes 7 critical biological activities or processes relevant to 1 or more of these 3 life stages and their outcomes, 15 habitat elements relevant to 1 or more of these 7 critical biological activities or processes for 1 or more life stages, and 9 controlling factors that affect 1 or more of these 15 habitat elements. Because the lower Colorado River (LCR) comprises a highly regulated system, the controlling factors exclusively concern human activities.

The seven critical biological activities and processes identified across all life stages are: disease, eating/foraging, molt, nest attendance, nest site selection, predation and brood parasitism, and temperature regulation. The 15 habitat elements identified across all life stages are: anthropogenic disturbance, brood size, canopy closure, community type, food availability, genetic diversity and infectious agents, local hydrology, parental feeding behavior, parental nest attendance, patch size, predator and cowbird density, soil salinity, tree size, understory density, and water availability. The nine controlling factors identified across all habitat elements are: fire management, grazing, irrigation, mechanical thinning, nuisance species introduction and management, pesticide/herbicide application, planting regime, recreational activities, and water storage-delivery system design and operation.

## RESULTS

The analysis of the causal relationships shows which critical biological activities and processes most strongly support or limit each life-stage outcome in the present system, which habitat elements most strongly affect the rates of these critical biological activities and processes, and which controlling factors most strongly affect the abundance, distribution, or condition of these habitat elements.

The analysis identifies several critical biological activities and processes that significantly affect survivorship across multiple life stages. Highlights of the results include the following:

- Disease, eating/foraging, and predation are the most important critical biological activities and processes affecting survival of all life stages. Depredation of nests can be high and has been shown to be the primary cause of nest failure among some species along the LCR (exceeding 80 percent) (Powell and Steidl 2000). The effects act at the landscape scale. Other processes such as molt and temperature regulation are important, but the effects are more indirect and less certain.
- Only two processes strongly and directly affect reproduction—nest attendance and nest site selection. These two critical biological activities and processes are especially important because they also affect nestling survival. Note, in this instance, reproduction refers to the laying of eggs; the survival of the nestlings is modeled separately in our process.

Finally, the analysis highlights several potentially important causal relationships about which scientific understanding remains low. These may warrant attention to determine if improved understanding might provide additional management options for improving VEFL survivorship and recruitment along the LCR. Specifically, the findings suggest a need to improve the understanding of:

- Eating/foraging and nest site selection are by far affected by the most habitat variables. They have been the most studied among the factors influencing this species, yet much uncertainty remains.
- Predation and brood parasitism is only strongly affected by two habitat elements—patch size and predator and cowbird density. Patch size affects predation rates because patch size affects the proportion of stand area near the edge of the patch, a factor which has been shown to influence predation (Chalfoun and Martin 2009; Theimer et al. 2011). Predator and cowbird density affects depredation rates (Lima 2009).

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

- Nest attendance is only strongly affected by brood size and predator and cowbird density. Brood size affects the amount of time adult VEFL must spend foraging versus attending the nest. Predator and cowbird density can influence nest defense behavior and thus nest attendance.
- Food availability and brood size drive the ability of VEFL in each life stage to acquire needed energy and nutrients.
- Disease and temperature regulation are important physiological concerns that can be impacted strongly by habitat elements such as canopy closure and local hydrology as well as the presence of genetic diversity and infectious agents.

The research questions and gaps in scientific knowledge identified in this modeling effort serve as examples of topics the larger scientific community could explore to improve the overall understanding of the ecology of VEFL. These questions may or may not be relevant to the goals of the LCR MSCP. As such, they are not to be considered guidance for Reclamation or the LCR MSCP, nor are these knowledge gaps expected to be addressed under the program.

# Chapter 1 – Introduction

This document presents a conceptual ecological model (CEM) for the vermilion flycatcher (*Pyrocephalus rubinus*) (VEFL). The purpose of this model is to help the Bureau of Reclamation (Reclamation), Lower Colorado River Multi-Species Conservation Program (LCR MSCP), identify areas of scientific uncertainty concerning VEFL ecology, the effects of specific stressors, the effects of specific management actions aimed at habitat and species restoration, and the methods used to measure VEFL habitat and population conditions. The CEM methodology follows that developed for the Sacramento-San Joaquin River Delta Regional Ecosystem Restoration Implementation Plan (DiGennaro et al. 2012), with modifications. (Note: Attachment 1 provides an introduction to the CEM process. We recommend that those unfamiliar with this process read the attachment before continuing with this document.)

The CEM addresses the VEFL population along the river and lakes of the lower Colorado River (LCR) and in other protected areas along the LCR managed as VEFL habitat. The model thus addresses the landscape as a whole rather than any single reach or managed area.

The most widely used sources of information for the VEFL conceptual ecological model are Rosenberg et al. (1991), Myers (2008), Ellison et al. (2009), Brand et al. (2010), and Raulston (2013). These publications summarize and cite large bodies of earlier studies. Where appropriate and accessible, those earlier studies are directly cited. The CEM also integrates numerous additional sources, particularly reports and articles completed since the aforementioned publications, information on current research projects, and the expert knowledge of LCR MSCP avian biologists. The purpose of the conceptual model is not to provide an updated literature review but to integrate the available information and knowledge into a CEM so it can be used for adaptive management.

This document is organized as follows: The remainder of chapter 1 provides a general description of the reproductive ecology of VEFL, the purpose of the model, and introduces the underlying concepts and structure of the CEM. Succeeding chapters present and explain the model for VEFL along the LCR and evaluate the implications of this information for management, monitoring, and research needs.

## **VERMILION FLYCATCHER REPRODUCTIVE ECOLOGY**

The VEFL is considered an incomplete migrant, with many populations remaining resident, but some northern populations migrating south for winter

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

(Ellison et al. 2009). Authors have suggested a combination of resident, breeding, and wintering birds occur within the LCR (Phillips et al. 1964; Rosenberg et al. 1991; Small 1994; Ellison et al. 2009). If so, wintering VEFL may depart the LCR by mid-April, whereas breeding birds may return to the LCR from their wintering grounds in late-February to mid-March to begin the breeding season (Rosenberg et al. 1991; Small 1994). Male birds arrive before females, with courtship beginning immediately upon the arrival of the females. Nest building commences soon after pairing, with the first egg laid shortly after nest completion (Rosenberg et al. 1991). Rosenberg et al. (1991) suggests that nesting sites are limited, causing heightened aggression between neighboring pairs that, in some cases, has delayed breeding.

VEFL nest sites are located in riparian woodlands, residential areas, and along the margins of agriculture with water nearby (Rosenberg et al. 1991; Great Basin Bird Observatory 2010). Rosenberg et al. (1991) suggested that nesting within the LCR was usually in cottonwoods or willows. However, recent studies have found VEFL use dry honey mesquite, similar to that in eastern and southern Arizona (Great Basin Bird Observatory 2010). The male VEFL suggests nests sites to the female during the courtship ritual (Ellison et al. 2009). The VEFL female builds the nest on a horizontal branch, usually in a fork. The typical clutch consists of two to three eggs, occasionally four, with two broods common during the breeding season (Rosenberg et al. 1991). Extra-pair copulation and intraspecific brood parasitism appear common (Ríos-Chelén et al. 2008). Brood parasitism by brown-headed cowbirds (*Molothrus ater*) ranges from 20 to 36 percent (Carothers 1974; Archer 1996; Ellison et al. 2007).

The onset of incubation is not well known, but all eggs usually hatch within 13 days of the final egg being laid (Ellison et al. 2009). Young birds fledge from the nest in 13–16 days, but it is unknown how long they are dependent upon their parents for food (Ellison et al. 2009).

Prey is captured using an “air sally” technique (Landres and MacMahon 1980). The diet of the VEFL has not been studied but likely includes bees, wasps, flies, beetles and grasshoppers (Rosenberg et al. 1991). VEFL have been observed feeding on fish on a few occasions (Andrews et al. 1996).

## **CONCEPTUAL ECOLOGICAL MODEL PURPOSES**

Adaptive management of natural resources requires a framework to help managers understand the state of knowledge about how a resource “works,” what elements of the resource they can affect through management, and how the resource will likely respond to management actions. The “resource” may be a population, species, habitat, or ecological complex. The best such frameworks

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

incorporate the combined knowledge of many professionals accumulated over years of investigations and management actions. CEMs capture and synthesize this knowledge (Fischenich 2008; DiGennaro et al. 2012).

CEMs explicitly identify: (1) the variables or attributes that best characterize resource conditions, (2) the factors that most strongly shape or control these variables under both natural and altered (including managed) conditions, (3) the character, strength, and predictability of the ways in which these factors do this shaping/controlling, and (4) how the characteristics of the resource vary as a result of the interplay of its shaping/controlling factors.

By integrating and explicitly organizing existing knowledge in this way, a CEM summarizes and documents: (1) what is known, with what certainty, and the sources of this information, (2) critical areas of uncertain or conflicting science that demand resolution to better guide management planning and action, (3) crucial attributes to use while monitoring system conditions and predicting the effects of experiments, management actions, and other potential agents of change, and (4) how the characteristics of the resource would likely change as a result of altering its shaping/controlling factors, including those resulting from management actions.

A CEM thus translates existing knowledge into a set of explicit hypotheses. The scientific community may consider some of these hypotheses well tested, but others less so. Through the model, scientists and managers can identify which hypotheses, and the assumptions they express, most strongly influence management actions. The CEM thus helps guide management actions based on the results of monitoring and experimentation. These results indicate whether expectations about the results of management actions – as clearly stated in the CEM – have been met or not. Both expected and unexpected results allow managers to update the model, improving certainty about some aspects of the model while requiring changes to other aspects, to guide the next cycle of management actions and research. The CEM, through its successive iterations, becomes the record of improving knowledge and the ability to manage the system.

## **CONCEPTUAL ECOLOGICAL MODEL STRUCTURE FOR THE VEFL**

The CEM methodology used here expands on that developed for the Sacramento-San Joaquin River Delta Regional Ecosystem Restoration Implementation Plan (DiGennaro et al. 2012). The expansion incorporates recommendations of Wildhaber et al. (2007, 2011), Kondolf et al. (2008), and Burke et al. (2009) to provide greater detail on causal linkages and outcomes and explicit demographic

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

notation in the characterization of life-stage outcomes (McDonald and Caswell 1993). Attachment 1 provides a detailed description of the methodology. The resulting model is a “life history” model, as is common for CEMs focused on individual species (Wildhaber et al. 2007, 2011). That is, it distinguishes the major life stages or events through which the individuals of a species must pass to complete a full life cycle, including reproducing, and the biologically crucial outcomes of each life stage. These biologically crucial outcomes typically include the number of individuals recruited to the next life stage (e.g., juvenile to adult) or next age class within a single life stage (recruitment rate), or the number of viable offspring produced (fertility rate). It then identifies the factors that shape the rates of these outcomes in the study area and thereby shapes the abundance, distribution, and persistence of the species in that area.

The VEFL conceptual ecological model has five core components as explained further in attachment 1:

- **Life stages** – These consist of the major growth stages and critical events through which the individuals of a species must pass in order to complete a full life cycle.
- **Life-stage outcomes** – These consist of the biologically crucial outcomes of each life stage, including the number of individuals recruited to the next life stage (e.g., juvenile to adult), or the number of viable eggs produced (fertility rate). The rates of the outcomes for an individual life stage depend on the rates of the critical biological activities and processes for that life stage.
- **Critical biological activities and processes** – These consist of the activities in which the species engages and the biological processes that take place during each life stage that significantly affect its life-stage outcomes rates. Examples of activities and processes for a bird species may include foraging, molt, nest site selection, and temperature regulation. Critical biological activities and processes typically are “rate” variables; the rate (intensity) of the activities and processes, taken together, determine the rate of recruitment of individuals to the next life stage.
- **Habitat elements** – These consist of the specific habitat conditions, the quality, abundance, and spatial and temporal distributions of which significantly affect the rates of the critical biological activities and processes for each life stage. These effects on critical biological activities and processes may be either beneficial or detrimental. Taken together, the suite of natural habitat elements for a life stage is called the “habitat template” for that life stage. Defining the natural habitat template may involve estimating specific thresholds or ranges of suitable values for particular habitat elements, outside of which one or more critical

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

biological activities or processes no longer fully support desired life-stage outcome rates – if the state of the science supports such estimates.

- **Controlling factors** – These consist of environmental conditions and dynamics – including human actions – that determine the quality, abundance, and spatial and temporal distributions of important habitat elements. Controlling factors are also called “drivers.” There may be a hierarchy of such factors affecting the system at different scales of time and space (Burke et al. 2009). For example, the availability of suitable nest sites for a riparian nesting bird may depend on factors such as canopy closure, humidity, and intermediate structure and community type, which in turn may depend on factors such as water storage-delivery system design and operation (dam design, reservoir morphology, and dam operations), which in turn is shaped by watershed geology, vegetation, climate, land use and water demand. The CEM identifies these five components and the causal relationships among them that affect life-stage outcome rates. Further, the CEM assesses each causal linkage based on four variables to the extent possible with the available information: (1) the character and direction of the effect, (2) the magnitude of the effect, (3) the predictability (consistency) of the effect, and (4) the status (certainty) of a present scientific understanding of the effect.

The CEM for each life stage thus identifies the causal relationships that most strongly support or limit the rates of its life-stage outcomes, support or limit the rate of each critical biological activity or process, and support or limit the quality, abundance, and distribution of each habitat element (as these affect other habitat elements or affect critical biological activities or processes). In addition, the model for each life stage highlights areas of scientific uncertainty concerning these causal relationships, the effects of specific management actions aimed at these relationships, and the suitability of the methods used to measure habitat and population conditions. Attachment 1 provides further details on the assessment of causal relationships, including the use of diagrams and a spreadsheet tool to record the details of the CEM and summarize the findings.

## Chapter 2 – VEFL Life Stage Model

A life stage consists of a biologically distinct portion of the life cycle of a species during which individuals undergo distinct developments in body form and function, engage in distinct behaviors, use distinct sets of habitats, and/or interact with their larger ecosystems in ways that differ from those associated with other life stages. This chapter proposes a life stage model for VEFL along the LCR on which to build the CEM.

### **INTRODUCTION TO THE VEFL LIFE CYCLE**

In the development of the CEM for VEFL, we could not find a complete demographic study of the species. We therefore chose to represent the VEFL with a three stage model to be consistent with other species documented within the LCR MSCP and to be most useful to management.

In many studies of avian demography, nest survival is considered integral in the reproduction of adults because adults are heavily invested in the care of eggs and nestlings (Etterson et al. 2011). We treat the nest stage as separate from adult reproduction due to the specific factors influencing the nest, the common creation of multiple broods by this species, and the fit with the life-stage outcome modelling structure used in this CEM process.

We have chosen to combine the egg and nestling phases of development into a nest stage because both the eggs and nestlings occupy the same nest; therefore, management focused on the nest will cover eggs and nestlings. Further, most research conducted on VEFL breeding has focused on the number of young fledged and not on the number of eggs hatched—meaning that most of the available information is on the habitat characteristics and management actions associated with success of the nest through both incubation and brooding periods.

The CEM is complicated by the lack of knowledge of the migratory behavior of the VEFL population occupying the LCR. Authors have suggested a combination of resident, breeding, and wintering birds occur within the LCR (Phillips et al. 1964; Rosenberg et al. 1991; Small 1994; Ellison et al. 2009). The conservation needs for the species within the LCR are likely highest for the breeding birds; thus, we focus the model on three life stages normally associated with the breeding season—nest, juvenile, and breeding adult.

## **VEFL LIFE STAGE 1 – NEST**

We consider the nest stage to be the first in the life cycle of the VEFL. It begins when the egg is laid and ends either when the young fledge or the nest fails. Eggs are usually laid in mid-March for a first brood, to as late as mid-June for a second brood (Ellison 2008; Ellison et al. 2009). Incubation lasts around 13 days, and young birds fledge from the nest in 13–16 days (Ellison et al. 2009). The life-stage outcome from the nest stage is the survival of eggs and associated nestlings until fledging. It is important to note that the outcome of the nest stage is inherently tied to the behavior and condition of the parents.

## **VEFL LIFE STAGE 2 – JUVENILE**

The juvenile stage begins at fledging and ends when the bird begins courtship activities the following spring. Juveniles are dependent upon the parents for some time after fledging, but the length of time has not been established (Ellison et al. 2009). For those birds migrating out of the area, they depart as early as late August (Small 1994). The life-stage outcome from the juvenile stage is the survival of the bird from fledging until the beginning of the courtship stage of the following breeding season. There are no studies available that analyze the survival rates of juveniles in this species; however, it may be assumed to be lower than adult survival rates that have been shown to be approximately  $0.48 \pm 0.21$  (Michel et al. 2006; Ricklefs and Shea 2007).

## **VEFL LIFE STAGE 3 – BREEDING ADULT**

The adult stage begins when the bird begins courtship activities in spring after its first winter. Generally, adults initiate courtship in late February or early March, with males arriving on the territory earlier if they left at all—and setting up territories before females arrive (Small 1994). Rosenberg et al. (1991) suggests that males staying on territory year round may breed earlier in spring with higher success.

The male VEFL suggests nest sites to the female during the courtship ritual (Ellison et al. 2009). The VEFL female builds the nest on a horizontal branch, usually in a fork. The typical clutch consists of two to three eggs, occasionally four, with two broods common during the breeding season (Rosenberg et al. 1991). Nest building commences soon after pairing, with the first egg laid shortly after nest completion (Rosenberg et al. 1991). Rosenberg et al. (1991) suggests that nesting sites are limited, causing heightened aggression between neighboring pairs that, in some cases, has delayed breeding up to a month.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

The life-stage outcomes for breeding adults are survival and reproduction—here defined as the production of eggs. Most studies of bird demography define fecundity—or the reproductive rates of adults—as the number of offspring fledged (Etterson et al. 2011). We have separated the nest stage from adult fecundity to more clearly display the information regarding nest success so that it can be better assessed by management. Therefore, adult reproduction involves the acts of pairing, site selection, nest building, and the production of eggs. Annual adult survival rates have been shown to be approximately  $0.48 \pm 0.21$  (Michel et al. 2006; Ricklefs and Shea 2007). The longevity record for VEFL, based on band recovery, is currently 5 years and 6 months (Klimkiewicz and Futcher 1989).

It is important to note that the post-breeding period is a significant part of a bird's life cycle. During the post-breeding period, adults may prospect for potential future breeding areas or move into habitat types that differ from breeding areas. VEFL are observed using more open areas, including residential areas, but are still in proximity to water during the non-breeding season (Rosenberg et al. 1991). Although male and female post-breeding individuals have different goals and responsibilities on the breeding grounds, we have included them all within the breeding adult life stage because their habitat use is similar, and thus, management directed at breeding adults will likely benefit all demographics present on the breeding grounds.


## LIFE STAGE MODEL SUMMARY

Based on this information, the VEFL conceptual ecological model distinguishes three life stages and their associated life-stage outcomes as shown in table 1 and figure 1. The life stages are numbered sequentially beginning with the nest.

Table 1.—VEFL life stages and outcomes in the LCR ecosystem

Life stage	Life-stage outcome(s)
1. Nest	<ul style="list-style-type: none"><li>• Survival</li></ul>
2. Juvenile	<ul style="list-style-type: none"><li>• Survival</li></ul>
3. Breeding adult	<ul style="list-style-type: none"><li>• Survival</li><li>• Reproduction</li></ul>

Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River



**Figure 1.—Proposed VEFL life history model.**

Squares indicate the life stages, and diamonds indicate the life-stage outcomes.

$S_{NJ}$  = survivorship rate, nest;  $S_{JB}$  = survivorship rate, juveniles;  $S_{BB}$  = survivorship rate, breeding adults; and  $R_{BN}$  = reproduction rate, breeding adults.

# Chapter 3 – Critical Biological Activities and Processes

Critical biological activities and processes consist of activities in which the species engages and biological processes that take place during each life stage that significantly shape the rate(s) of the outcome(s) for that life stage. Critical biological activities and processes are “rate” variables (i.e., the rate [intensity] of these activities and processes, taken together, determine the rate of recruitment of individuals from one life stage to the next).

The CEM identifies seven critical biological activities and processes that affect one or more VEFL life stages. Some of these activities or processes differ in their details among life stages. However, grouping activities or processes across all life stages into broad types makes it easier to compare the individual life stages to each other across the entire life cycle. Table 2 lists the seven critical biological activities and processes and their distribution across life stages.

Table 2.—Distribution of VEFL critical biological activities and processes among life stages

(Xs indicate that the critical biological activity or process is applicable to that life stage.)

Life stage →			
	Nest	Juvenile	Breeding adult
Critical biological activity or process ↓			
Disease	X	X	X
Eating/foraging	X	X	X
Molt	X	X	X
Nest attendance			X
Nest site selection			X
Predation and brood parasitism	X	X	X
Temperature regulation	X	X	X

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

The most widely used sources of the information used to identify the critical biological activities and processes are Rosenberg et al. (1991), Myers (2008), Ellison et al. (2009), Brand et al. (2010), and Raulston (2013). The identification also integrates information from both older and more recent works as well as the expert knowledge of LCR MSCP biologists. The following paragraphs discuss the seven critical biological activities and processes in alphabetical order.

## **DISEASE**

This process refers to diseases caused either by lack of genetic diversity or by infectious agents, including the effects of endo- and ecto-parasites. Little research has focused on specific diseases inflicting the VEFL. However, there is a wealth of knowledge regarding avian diseases and parasites that affect passerine birds within North America, which indicates a large number of diseases (Morishita et al. 1999) can be difficult to detect (Jarvi et al. 2002) and can have differing effects on different species (Merino et al. 2000; Palinauskas et al. 2008). VEFL at all life stages are conceivably susceptible to disease.

## **EATING/FORAGING**

This process applies to all life stages but somewhat differently. Nestlings are fed by adults and do not forage. Juveniles forage on their own but are still fed by adults shortly after fledging, and this feeding may be critical to their survival. The ability of nestlings and juveniles to eat is determined by the provisioning rate of its parents and competition with brood mates. Breeding adults must forage to feed themselves and their dependent young. Prey is captured using an “air sally” technique (Landres and MacMahon 1980). The diet of the VEFL has not been studied but likely includes bees, wasps, flies, beetles, and grasshoppers (Rosenberg et al. 1991). VEFL have been observed feeding on fish on a few occasions (Andrews et al. 1996). Foraging is done by juveniles and adults, but it is important to note that foraging by the parents affects the provisioning rate to nestlings and nest attendance by adults.

## **MOLT**

The VEFL uses a complex alternate molt strategy (Ellison et al. 2009; Howell 2010). Nestling VEFL must molt from natal down into juvenal plumage while in the nest. The success of this molt is dependent upon the adult provisioning rate. Molting is an energetically costly process that may make nestlings more susceptible to death when resources are scarce. Feather quality may be negatively affected by poor diet, and the nestlings may compensate by shifting resources

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

from other critical functions such as the immune system, putting them at further risk (Birkhead et al. 1999). Similarly, adult birds molt after the breeding season and before autumn migration (if not resident) and face the same challenges as nestlings (Howell 2010).

## **NEST ATTENDANCE**

The female VEFL does all of the incubating and brooding (Taylor and Hanson 1970; Ellison et al. 2009). After hatching, both parents feed the young and the male occasionally feeds the brooding female (Taylor and Hanson 1970). During the days after hatching, females have been observed attending the nest 60–80 percent of the time (Taylor and Hanson 1970).

## **NEST SITE SELECTION**

Both breeding males and females select a nest site, with males selecting territories and females selecting the actual nest site within that territory (Taylor and Hanson 1970; Ellison et al. 2009). Nest site selection is important for reproductive success because nest success varies spatially as a result of vegetation characteristics, food availability, predator types and densities, hydrology, and unique events such as flooding (Powell and Steidl 2000; Lima 2009; Ellison et al. 2009; Brand et al. 2010).

## **PREDATION AND BROOD PARASITISM**

Nest predation and brood parasitism certainly affect the success of songbird nests within the LCR (Powell and Steidl 2000; Brand et al. 2010). Powell and Steidl (2000) found that nest predation was more significant than parasitism, accounting for 81 percent of nest failures. Brood parasitism by brown-headed cowbirds ranges from 20 percent to 36 percent among successful nests (Carothers 1974; Archer 1996; Ellison et al. 2007). Nest predator and cowbird abundance is influenced by patch size and the nest's relative proximity to a vegetation edge (Rosenberg et al. 1999; Winfree 2004). These two processes have been combined for the nestling and egg stages because (1) cowbirds are both nest predators and brood parasites (Theimer et al. 2011) and (2) habitat characteristics (distance to edge, patch width, etc.) affect both processes similarly (Rosenberg et al. 1999).

Juvenile and adult birds are also vulnerable to predation, but it has not been studied in VEFL. Predation on juveniles and adults is not as easily quantified but affects juveniles and adults and indirectly affects nestling survival through nest

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**

abandonment after predation of a breeding adult bird. Predation risk can result in many behavioral adaptations in passerines, including nest locations, densities, clutch size, egg size, etc. (Lima 2009).

## **TEMPERATURE REGULATION**

Temperature regulation is important for any organism inhabiting a region with temperatures as high as that along the LCR. Although overheating is possible during all life stages, most of the concern has been toward eggs and nestlings (Hunter et al. 1987a, 1987b; Rosenberg et al. 1991). Adults can affect the temperature regulation of eggs and nestlings through their own behavior (incubation, brooding, or shading) and through nest placement.

# Chapter 4 – Habitat Elements

Habitat elements consist of specific habitat conditions that ensure, allow, or interfere with critical biological activities and processes.

This chapter identifies 15 habitat elements that affect 1 or more critical biological activities or processes across the 3 VEFL life stages. Some of these habitat elements differ in their details among life stages. For example, VEFL at different life stages experience different predation. However, using the same labels for the same *kinds* of habitat elements across all life stages makes comparison and integration of the CEMs for the individual life stages across the entire life cycle less difficult. Table 3 lists the 15 habitat elements and the critical biological activities and processes that they *directly* affect across all VEFL life stages.

Table 3.—Distribution of VEFL habitat elements and the critical biological activities and processes that they directly affect across all life stages (Xs indicate that the habitat element is applicable to that critical biological activity or process.)

Critical biological activity or process →							
Habitat element ↓	Disease	Eating/foraging	Molt	Nest attendance	Nest site selection	Predation and brood parasitism	Temperature regulation
Anthropogenic disturbance		X		X	X	X	
Brood size		X		X			
Canopy closure		X			X	X	X
Community type					X	X	
Food availability		X		X	X		
Genetic diversity and infectious agents	X						
Local hydrology					X		X
Parental feeding behavior		X					
Parental nest attendance		X				X	X
Patch size					X	X	
Predator and cowbird density		X		X	X	X	
Soil salinity							
Tree size					X		
Understory density					X		
Water availability					X		X

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

The diagrams and other references to habitat elements elsewhere in this document identify the habitat elements by a one-to-three-word short name. However, each short name in fact refers to a longer, complete name. For example, “predator and cowbird density” is the short name for “The abundance and distribution of predators and brood parasites.” The following paragraphs provide the full name for each habitat element and a detailed definition, addressing the elements in alphabetical order.

The most widely used sources of the information used to identify the habitat elements are Rosenberg et al. (1991), Myers (2008), Ellison et al. (2009), Brand et al. (2010), and Raulston (2013). The identification also integrates information from both older and more recent works as well as the expert knowledge of LCR MSCP biologists.

VEFL nest sites are located in riparian woodlands, residential areas, and along the margin of agriculture with water nearby (Rosenberg et al. 1991). VEFL in the Southwest United States tend to nest at higher densities in native cottonwoods rather than in other trees (Brand et al. 2010). The use of dry honey mesquite has been observed along the LCR but only in created park-like settings (Great Basin Bird Observatory 2010).

Prey is captured using an “air sally” technique (Landres and MacMahon 1980). The diet of the VEFL has not been studied but likely includes bees, wasps, flies, beetles, and grasshoppers (Rosenberg et al. 1991). VEFL have been observed feeding on fish on a few occasions (Andrews et al. 1996).

As with all tabulations of habitat associations, inferences that particular habitat characteristics are critical to a species or life stage require evidence and CEMs for why each association matters to species viability (Rosenfeld 2003; Rosenfeld and Hatfield 2006.)

## **ANTHROPOGENIC DISTURBANCE**

*Full name:* **Human activity within or surrounding a given habitat patch, including noise, pollution, and other disturbances associated with human activity.** Anthropogenic disturbance can affect both breeding success and survival of birds (reviewed by Barber et al. 2010; Francis and Barber 2013). Noise might mask conspecific cues such as songs or calls—making it more difficult for VEFL to attract or find mates or defend territories. Noise can shift the foraging/vigilance tradeoff – either putting an individual at higher risk due to starvation or to predation (Ware et al. 2015). Noise can cause behavioral changes, physiological changes, and species diversity changes within an area. The VEFL has been observed to sing longer songs in noisier environments (Ríos-Chelén et al. 2012).

## BROOD SIZE

*Full name:* **The number of young in the nest.** This element refers to the number of young that the parents must rear. It differs from clutch size, which refers to the number of eggs laid. Brood size is related to maternal health, and the well-being of both parents depends in part on the availability of sufficient food resources in close proximity to the breeding territory as well as other factors such as predator and cowbird density (see “Predator and Cowbird Density,” below). A VEFL typical clutch consists of two to three eggs, occasionally four, with two broods common during the breeding season (Rosenberg et al. 1991).

## CANOPY CLOSURE

*Full name:* **The proportion of the sky hemisphere obscured by vegetation when viewed from a single point as measured with a spherical densitometer (Jennings et al. 1999).** This element refers to the percent canopy closure of canopy vegetation in the vicinity of the VEFL nest site. Canopy closure of riparian vegetation has been shown to be important to VEFL, but higher canopy densities appear to be avoided (Ellison et al. 2009). Studies conducted within the LCR have not found a predictive formula for VEFL occupancy (Raulston 2013).

## COMMUNITY TYPE

*Full name:* **The species composition of the riparian forest patch.** This element refers to the species composition of riparian habitat used for breeding by VEFL. Research shows that flycatchers are adaptable, able to use various types of native and non-native broadleaf deciduous habitats at different elevations (Brand et al. 2010). However, VEFL densities are substantially higher in native habitats such as large cottonwood stands within the LCR region (Brand et al. 2010).

## FOOD AVAILABILITY

*Full name:* **The abundance of food available for adults and their young.** This element refers to the taxonomic and size composition of the invertebrates that an individual VEFL will encounter during each life stage as well as the density and spatial distribution of the food supply in proximity to the nest. The diet of the VEFL has not been studied but likely includes bees, wasps, flies, beetles, and grasshoppers (Rosenberg et al. 1991). VEFL have been observed feeding on fish on a few occasions (Andrews et al. 1996). The abundance and condition of

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

the food supply affects adult health, growth and development of nestlings and juveniles, the progress of molt, and the success of later stages in the annual cycle (i.e., migration).

## **GENETIC DIVERSITY AND INFECTIOUS AGENTS**

*Full name:* **The genetic diversity of VEFL individuals and the types, abundance, and distribution of infectious agents and their vectors.** The genetic diversity component of this element refers to the genetic homogeneity versus heterogeneity of a population during each life stage. The greater the heterogeneity, the greater the possibility that individuals of a given life stage will have genetically encoded abilities to survive their encounters with the diverse stresses presented by their environment and/or take advantage of the opportunities presented. The genetic diversity of the LCR populations of VEFL has not been studied.

The infectious agent component of this element refers to the spectrum of viruses, bacteria, fungi, ecto-parasites, and endo-parasites that individual VEFL are likely to encounter during each life stage. There have been no specific studies of the infectious agents and their effects on the VEFL within the LCR. However, there is a wealth of knowledge regarding avian diseases and parasites that affect passerine birds within North America, which indicates a large number of diseases (e.g., Morishita et al. 1999) can be difficult to detect (Jarvi et al. 2002) and can have differing effects on different species (Merino et al. 2000; Palinauskas et al. 2008).

## **LOCAL HYDROLOGY**

*Full name:* **Aspects such as the distance to standing water or the presence of adjacent water bodies, the timing and volume of floods, depth to the water table, and soil moisture levels.** This element refers to anything that affects soil moisture, such as the proximity of water to the nesting habitat, elevation, irrigation practices, and soil texture. The local hydrological conditions affect other aspects of habitat such as vegetation structure and abundance of arthropods. Wetter conditions might also provide cooler temperatures and more humid conditions necessary for egg and chick survival in desert systems (Rosenberg et al. 1991).

## PARENTAL FEEDING BEHAVIOR

*Full name:* **The ability and behavior of parents to feed and care for nestlings and juveniles after they fledge from the nest.** This element refers to the capacity of both parents to provision food for nestlings and recently fledged birds. This rate is dependent upon food availability and the number of young in the brood. This rate influences the amount of food and time spent foraging by juvenile birds.

## PARENTAL NEST ATTENDANCE

*Full name:* **The ability of both parents to care for young during the egg/incubation and nestling stages.** This element refers to the capacity of both parents to share nesting and brood rearing responsibilities until fledging. It is affected by the presence of predators and competitors, food availability, and the ability to thermal regulate. Female VEFL have the sole responsibility of brooding young, but the territory is actively defended by the male (Ellison et al. 2009).

## PATCH SIZE

*Full name:* **The size of riparian habitat patches.** This element refers to the areal extent of a given patch of riparian vegetation. Patch size affects the number of breeding pairs that an area can support as well as the density of predators, competitors, and brood parasites. Little data are available regarding the importance of patch size on VEFL breeding success.

## PREDATOR AND COWBIRD DENSITY

*Full name:* **The abundance and distribution of predators and brood parasites.** This element refers to a set of closely related variables that affect the likelihood that different kinds of predators will encounter and successfully prey on VEFL during the egg, nestling, or adult life stages or that cowbirds or other nest parasites will lay eggs in the nest. The variables of this element include the species and size of the fauna that prey on VEFL during different life stages, the density and spatial distribution of these fauna in the riparian habitat used by VEFL, and whether predator activity may vary in relation to other factors (e.g., time of day, patch size and width, matrix community type, etc.). Powell and Steidl (2000) report observing that 81 percent of the southwestern riparian nests they studied were predated with a large amount of brood parasitism as well. Ellison et al. (2009) report that there were few records of brood parasitism by

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

cowbirds prior to 1974. However, a number of more recent reports suggest that parasitism is common and that VEFL have little natural defense (Carothers 1974; Mason 1986; Ellison et al. 2007). Brood parasitism by brown-headed cowbirds ranges from 20 to 36 percent among successful nests (Carothers 1974; Archer 1996; Ellison et al. 2007). The effect of predator and cowbird density can have impacts more subtle than survival by altering prey behavior, nest site selection, breeding behavior, and foraging behavior (Lima 1998, 2009; Chalfoun and Martin 2009).

## **SOIL SALINITY**

*Full name:* **The salt content within the root zone of the soil (0–30 inches) as measured by electrical conductivity of the saturation extract value in decisiemens per meter at 25 degrees Celsius** (San Joaquin River Restoration Program 2014). Soil salinity can impact the vigor of various plant species to different degrees and can ultimately influence plant community type and structure (San Joaquin River Restoration Program 2014).

## **TREE SIZE**

*Full name:* **The diameter of a tree at breast height, averaged across the stand.** VEFL are known to nest in large trees within the LCR, most often cottonwood, but also honey mesquite in lower densities (Brand et al. 2010; Great Basin Bird Observatory 2010).

## **UNDERSTORY DENSITY**

*Full name:* **The density of the understory layer of vegetation (0-1.5 meters) (Powell and Steidl 2002).** VEFL appear to select for low understory density, including lower vegetative coverage, lower volume, and fewer shrubs (Great Basin Bird Observatory 2010).

## **WATER AVAILABILITY**

*Full name:* **The availability of nearby surface water.** The presence of breeding VEFL has been highly correlated with the presence of nearby surface water, either natural or artificial (Great Basin Bird Observatory 2010).

## Chapter 5 – Controlling Factors

Controlling factors consist of environmental conditions and dynamics, both natural and anthropogenic, which significantly affect the abundance, spatial and temporal distributions, and quality of critical habitat elements. These may also significantly and directly affect some critical biological activities or processes. A hierarchy of such factors exists, with long-term dynamics of climate and geology at the top. However, this CEM focuses on nine immediate controlling factors that are within the scope of potential human manipulation. The nine controlling factors identified in this CEM do not constitute individual variables; rather, each identifies a category of variables (including human activities) that share specific features that make it useful to treat them together. Table 4 lists the nine controlling factors and the habitat elements they directly affect. Table 4 shows two habitat elements that are not directly affected by any controlling factor (brood size and genetic diversity and infectious agents). These latter habitat elements are directly shaped by the condition of one or more other habitat elements rather than by any of the controlling factors.

Table 4.—Habitat elements directly affected by controlling factors

Controlling factor →	Fire management	Grazing	Irrigation	Mechanical thinning	Nuisance species introduction and management	Pesticide/herbicide application	Planting regime	Recreational activities	Water storage-delivery system design and operation
Habitat element ↓	Fire management	Grazing	Irrigation	Mechanical thinning	Nuisance species introduction and management	Pesticide/herbicide application	Planting regime	Recreational activities	Water storage-delivery system design and operation
Anthropogenic disturbance		X		X		X	X	X	
Brood size	N/A*								
Canopy closure	X			X	X		X	X	
Community type	X	X			X		X	X	X
Food availability					X	X	X		
Genetic diversity and infectious agents	N/A*								
Local hydrology			X						X
Parental feeding behavior								X	
Parental nest attendance								X	
Patch size	X	X		X	X		X	X	
Predator and cowbird density					X		X	X	
Soil salinity			X						X
Tree size	X			X	X		X		
Understory density	X	X		X	X		X	X	X
Water availability									X

\* N/A values suggest that none of the identified controlling factors *directly* affect the habitat element.

## **FIRE MANAGEMENT**

This factor addresses any fire management (whether prescribed fire or fire suppression) that could affect VEFL or their habitat. Effects may include creation of habitat that supports or excludes VEFL, a reduction in the food supply of invertebrates, or support of species that pose threats to VEFL as predators, competitors, or carriers of infectious agents. While fire can decrease understory vegetation to the benefit of VEFL, most of the other impacts are negative, such as decreased canopy closure, decreased patch size, decreased tree size, and increased soil temperature, which in turn decreases the food supply.

Climate change is also projected to affect fire frequency along the LCR (U.S. Fish and Wildlife Service [USFWS] 2013).

## **GRAZING**

This factor addresses the grazing activity on riparian habitats along the LCR and in surrounding areas that could affect VEFL or their habitat. Grazing by cattle, burros, or mule deer across the arid Southwestern United States has substantially degraded riparian habitat (see USDA Forest Service 1979; Rickard and Cushing 1982; Cannon and Knopf 1984; Klebenow and Oakleaf 1984; General Accounting Office 1988; Clary and Webster 1989; Schultz and Leininger 1990; and Belsky et al. 1999 in USFWS 2002). (Note: Reclamation staff and researchers have observed mule deer browsing on LCR sites, which may become an issue if populations are not managed). Grazing may thin the understory, which could benefit VEFL in the short term, but can also prevent the establishment of cottonwood, willow, or mesquite seedlings (Kauffman et al. 1997; Powell and Steidl 2002). VEFL occur in lower densities in grazed areas but respond quickly after grazing is stopped (Krueper et al. 2003).

## **IRRIGATION**

This factor addresses the human activities of artificially introducing water to the landscape to influence habitat. In many cases, irrigation may be implemented to simulate more natural riparian processes or to manage soil salinity levels.

## **MECHANICAL THINNING**

This factor addresses the active removal of vegetation from areas within the LCR region. Effects may include the creation of habitat that supports or excludes

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

VEFL or support of species that pose threats to VEFL such as predators, competitors, or carriers of infectious agents. This factor includes the thinning of vegetation within both riparian and matrix communities.

## **NUISANCE SPECIES INTRODUCTION AND MANAGEMENT**

This factor addresses the intentional or unintentional introduction of nuisance species (animals and plants) and their control that affects VEFL survival and reproduction. Nuisance species may infect, prey on, compete with, or present alternative food resources for VEFL during one or more life stages; cause other alterations to the riparian food web that affect VEFL; or affect physical habitat features such as canopy or shrub cover. For example, although VEFL successfully nest in sites dominated by invasive tamarisk (*Tamarix* spp.), they do so in lower densities, and tamarisk may negatively affect habitat in other ways (e.g., by lowering the water table) (Di Tomaso 1998; Brand et al. 2010).

The complicated nature of the relationship between tamarisk and VEFL is highlighted by another introduced species—the tamarisk beetle (*Diorhabda carinulata*). The tamarisk beetle was introduced to the LCR region in order to control invasive tamarisk (Bateman et al. 2013). However, defoliation of tamarisk due to beetle infestation causes decreases in humidity and cover along with increases in temperature (Bateman et al. 2013), thereby potentially degrading areas dominated by tamarisk as habitat for VEFL.

## **PESTICIDE/HERBICIDE APPLICATION**

This factor addresses pesticide/herbicide applications that may occur on or adjacent to riparian habitat of the LCR region. Effects may include sublethal poisoning of VEFL via ingestion of treated insects, pollution of runoff into wetland habitats that are toxic to prey of VEFL, and a reduced invertebrate food supply.

## **PLANTING REGIME**

This factor addresses the active program to restore cottonwood-willow riparian habitat along the LCR and includes both the community planted as well as the manner in which it is planted within restoration areas (e.g., density, age, and patch size). The composition of the species planted can affect not only the vertical and horizontal structure of the vegetation but also the insect community within a given patch.

## **RECREATIONAL ACTIVITIES**

This factor addresses the disturbance to VEFL from recreational activities. Even non-consumptive human activity can have negative effects on wildlife (reviewed by Boyle and Samson 1985). This is a broad category that encompasses the types of activity (e.g., boating, fishing, horseback riding, camping, etc.) as well as the frequency and intensity of those activities. The impacts may consist of disturbance and habitat alteration.

## **WATER STORAGE-DELIVERY SYSTEM DESIGN AND OPERATION**

Much of the habitat currently used by VEFL within the LCR is along regulated waterways. The water moving through this system is highly regulated for storage and delivery (diversion) to numerous international, Federal, State, Tribal, and municipal users and for hydropower generation.

The dynamic nature of a free-flowing river creates a mosaic of riparian habitats, and thus, a natural flow regime might be beneficial to VEFL. Natural floods can decrease understory vegetation or increase the presence of surface water, improving VEFL habitat (Great Basin Bird Observatory 2010).

## Chapter 6 – Conceptual Ecological Model by Life Stage

This chapter contains three sections, each presenting the CEM for a single VEFL life stage. The text and diagrams identify the critical biological activities and processes for each life stage, the habitat elements that support or limit the success of these critical biological activities and processes, the controlling factors that determine the abundance and quality of these habitat elements, and the causal links among them. The CEM sections specifically refer to the river and lakes of the LCR and other protected areas managed as VEFL habitat and thus address this landscape as a whole rather than any single reach or managed area.

The CEM for each life stage assesses the character and direction, magnitude, predictability, and scientific understanding of each causal link based on the following definitions (see attachment 1 for further details):

- **Character and direction** categorizes a causal relationship as positive, negative, or complex. “Positive” means that an increase in the causal node results in an increase in the affected node, while a decrease in the causal node results in a decrease in the affected node. “Negative” means that an increase in the causal node results in a decrease in the affected element, while a decrease in the causal node results in an increase in the affected node. Thus, “positive” or “negative” here do *not* mean that a relationship is beneficial or detrimental. The terms instead provide information analogous to the sign of a correlation coefficient. “Complex” means that there is more going on than a simple positive or negative relationship. Positive and negative relationships are further categorized based on whether they involve any response threshold in which the causal agent must cross some value before producing an effect. In addition, the “character and direction” attribute categorizes a causal relationship as uni- or bi-directional. Bi-directional relationships involve a reciprocal relationship in which each node affects the other.
- **Magnitude** refers to “...the degree to which a linkage controls the outcome *relative to other drivers*” (DiGennaro et al. 2012). Magnitude takes into account the spatial and temporal scale of the causal relationship as well as the strength (intensity) of the relationship at any single place and time. The present methodology separately rates the intensity, spatial scale, and temporal scale of each link on a three-part scale from “Low” to “High” and assesses overall link magnitude by averaging the ratings for these three. If it is not possible to estimate the intensity, spatial scale, or temporal scale of a link, the subattribute is rated as “Unknown” and ignored in the averaging. If all three subattributes are “Unknown,” however, the overall link magnitude is rated as “Unknown.” Just as the

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**


terms for link character provide information analogous to the sign of a correlation coefficient, the terms for link magnitude provide information analogous to the size of a correlation coefficient.

- **Predictability** refers to “...the degree to which current understanding of the system can be used to predict the role of the driver in influencing the outcome. Predictability ... captures variability... [and recognizes that] effects may vary so much that properly measuring and statistically characterizing inputs to the model are difficult” (DiGennaro et al. 2012). A causal relationship may be unpredictable because of natural variability in the system or because its effects depend on the interaction of other factors with independent sources for their own variability. Just as the terms for link character provide information analogous to the sign of a correlation coefficient, the terms for link predictability provide information analogous to the size of the range of error for a correlation coefficient. The present methodology rates the predictability of each link on a three-part scale from “Low” to “High.” If it is not possible to rate predictability due to a lack of information, then the link is given a rating of “Unknown” for predictability.
- **Scientific understanding** refers to the degree of agreement represented in the scientific literature and among experts in understanding how each causal relationship works—its character, magnitude, and predictability. Link predictability and understanding are independent attributes. A link may be highly predictable but poorly understood or poorly predictable but well understood. The present methodology rates the state of scientific understanding of each link on a three-part scale from “Low” to “High.”

**The CEM for each life stage thus identifies the causal relationships that most strongly support or limit life-stage outcomes, support or limit the rate of each critical biological activity or process, and support or limit the quality of each habitat element, as that element affects other habitat elements or affects critical biological activities or processes.**

A separate spreadsheet is used to record the assessment of the character and direction, magnitude, predictability, and scientific understanding for each causal link along with the underlying rationale and citations for each life stage. The CEM for each life stage, as cataloged in its spreadsheet, is illustrated with diagrams showing the controlling factors, habitat elements, critical biological activities and processes, and causal links identified for that life stage. A diagram may also visually display information on the character and direction, magnitude, predictability, and/or scientific understanding of every link. The diagrams use a common set of conventions for identifying the controlling factors, habitat elements, critical biological activities and processes, and life-stage outcomes as well as for displaying information about the causal links. Figure 2 illustrates these conventions.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**



**Figure 2.—Diagram conventions for LCR MSCP conceptual ecological models.**

The discussion of each life stage includes an analysis of the information contained in the spreadsheet. The analyses highlight causal chains that strongly affect survivorship, identify important causal relationships with different levels of predictability, and identify important causal relationships with high scientific uncertainty. The latter constitutes topics of potential importance for adaptive management investigation.

The causal relationships between controlling factors and habitat elements are essentially identical across all three life stages. For this reason, the discussion of controlling factor-habitat element linkages across all three life stages appears in a subsequent chapter.

## VEFL LIFE STAGE 1 – NEST

The nest stage lasts from when the egg is laid until either the young fledge or the nest fails. Success during this life stage – successful transition to the juvenile stage – involves organism survival, maturation, molt, and fledging. The organisms actively interact with their environment.

The CEM (figures 3 and 4) recognizes five (of seven) critical biological activities and processes for this life stage, and they are presented here as they appear on the following figures:

1. **Disease** – Although the literature does not emphasize disease as affecting population levels of VEFL, we still feel that disease bears mentioning. Diseases and parasites are prevalent in avian populations, so it is safe to assume they have an impact on VEFL (Morishita et al. 1999,; Lachish et al. 2011). Disease and parasite impacts along the LCR are areas recommended for further research.

The CEM recognizes genetic diversity and infectious agents as a habitat element affecting disease.

2. **Eating/Foraging** – The nestling must eat to maintain metabolic processes.

The CEM recognizes anthropogenic disturbance, brood size, parental feeding behavior, and parental nest attendance as habitat elements affecting eating.

3. **Molt** – The nestling must molt into juvenile plumage.

The CEM does not recognize any habitat elements as directly affecting molt.

Other critical activities influencing molt indirectly include those affecting energy resources such as disease and eating.

4. **Predation and Brood Parasitism** – Both nest predation and brood parasitism affect the survival of a nest and are affected by similar habitat elements. Brood parasitism has been identified as a threat to VEFL. High rates of predation and brood parasitism have been observed for VEFL and for other species within the LCR, accounting for over 80 percent of nest failures in some species (Powell and Steidl 2000). Rasmussen and Sealy (2006) further speculate that the impact of brood parasitism post-fledging may be much higher than assumed.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

The CEM recognizes canopy closure, community type, parental nest attendance, patch size, and predator and cowbird density as habitat elements affecting predation and brood parasitism. Note that the effects of anthropogenic disturbance on this element are not well known.

5. **Temperature Regulation** – The eggs and nestlings must maintain an optimum temperature to develop and survive.

The CEM recognizes canopy closure, disease, local hydrology, parental nest attendance, and water availability as the primary habitat elements directly affecting temperature regulation.

Disease and eating are two critical biological activities that can influence temperature regulation.




Figure 3.—VEFL life stage 1 – nest, basic CEM diagram. Only elements with connections within this life stage are presented.




Figure 4.—VEFL life stage 1 – nest, high- and medium-magnitude relationships. Only elements with high- or medium-magnitude connections within this life stage are presented.

## VEFL LIFE STAGE 2 – JUVENILE

The juvenile stage begins at fledging and ends when the bird engages in breeding activities, usually the following year. Success during this life stage – successful transition to the next stage – involves organism survival and maturation. The organisms actively interact with their environment.

The CEM (figures 5 and 6) recognizes five (of seven) critical biological activities and processes for this life stage, and they are presented here as they appear on the following figures:

1. **Disease** – Although the literature does not emphasize disease as affecting population levels of VEFL, we still feel that disease bears mentioning. Diseases and parasites are prevalent in avian populations, so it is safe to assume they have an impact on VEFL (Morishita et al. 1999; Lachish et al. 2011). Disease and parasite impacts along the LCR are areas recommended for further research.

The CEM recognizes genetic diversity and infectious agents as a habitat element affecting disease.

2. **Eating/Foraging** – The juvenile must eat to maintain metabolic processes. While they will increasingly forage on their own, they are dependent upon food from their parents for an unknown period of time after fledging.

The CEM recognizes anthropogenic disturbance, brood size, canopy closure, food availability, parental feeding behavior, and predator and cowbird density as habitat elements affecting foraging. In addition, disease can affect the foraging efficiency of a juvenile.

3. **Molt** – The juvenile must molt into adult plumage.

The CEM does not recognize any habitat elements as directly affecting molt.

Other critical activities influencing molt include those affecting energy resources such as disease, eating, and foraging.

4. **Predation and Brood Parasitism** – The juvenile is at risk for direct predation and may still suffer from previous brood parasitism. Rasmussen and Sealy (2006) speculate that the impact of brood parasitism post-fledging may be much higher than assumed, as adult birds continue to feed the parasite nestlings.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**

The CEM recognizes anthropogenic disturbance, canopy closure, community type, patch size, and predator and cowbird density as habitat elements affecting predation.

5. **Temperature Regulation** – The juvenile must maintain an optimum temperature to survive.

The CEM recognizes canopy closure, local hydrology, and water availability as habitat elements directly affecting temperature regulation. Disease is a critical process that can also influence temperature regulation.




Figure 5.—VEFL life stage 2 – juvenile, basic CEM diagram. Only elements with connections within this life stage are presented.




Figure 6.—VEFL life stage 2 – juvenile, high- and medium-magnitude relationships. Only elements with high- or medium-magnitude connections within this life stage are presented.

## VEFL LIFE STAGE 3 – BREEDING ADULT

The breeding adult stage begins when the bird engages in breeding activities after its first or subsequent winter. Success during this life stage – measured on an annual basis – involves organism survival and breeding. Individuals that do not successfully find a territory, floaters, are also included in this category even though they do not breed. The organisms actively interact with their environment.

The CEM (figures 7 and 8) recognizes seven (of seven) critical biological activities and processes for this life stage, and they are presented here as they appear on the following figures:

1. **Disease** – Although the literature does not emphasize disease as affecting population levels of VEFL, we still feel that disease bears mentioning. Diseases and parasites are prevalent in avian populations, so it is safe to assume they have an impact on VEFL (Morishita et al. 1999; Lachish et al. 2011). Disease and parasite impacts along the LCR are areas recommended for potential research.

The CEM recognizes genetic diversity and infectious agents as a habitat element affecting disease.

2. **Eating/Foraging** – The breeding adult must forage to feed itself and its young. Both their survival and the survival of their young are dependent upon the foraging rate, which can be influenced by a number of factors.

The CEM recognizes anthropogenic disturbance, brood size, canopy closure, food availability, parental feeding behavior, parental nest attendance, and predator and cowbird density as primary habitat elements affecting foraging. Disease may cause a decrease in foraging performance.

3. **Molt** – The adult must molt annually to maintain flight capabilities.

The CEM does not recognize any habitat elements as directly affecting molt.

Other critical activities influencing molt include those affecting energy resources such as disease and foraging.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**

4. **Nest Attendance** – The breeding adult must attend the nest to incubate eggs, brood young, and feed young. This primary responsibility falls upon the female, but the male does provide food and nest defense.

The CEM recognizes anthropogenic disturbance, brood size, food availability, and predator and cowbird density as the major factors affecting nest attendance. Disease may have an effect, but it remains less certain.

5. **Nest Site Selection** – This process includes both territory establishment and the placement of nests. Territory establishment is especially important because if a bird fails to establish a territory (or find a male with a territory in the case of females), the bird will be a floater and is unlikely to breed during that season. The breeding adult must choose where to place territories and nests, thereby affecting breeding success.

The CEM recognizes canopy closure, community type, and patch size as primary habitat elements affecting nest site selection. In addition, local hydrology and predator and cowbird density may act as secondary habitat elements influencing nest site selection.

6. **Predation** – VEFL adults are conspicuous, and their perching and sallying foraging behavior makes them susceptible to depredation. Adults must avoid predators to survive.

The CEM recognizes patch size and predator and cowbird density as the primary elements affecting predation. Canopy closure may also have influences. Note that the effects of anthropogenic disturbance on this element are not well known.

7. **Temperature Regulation** – The adult must maintain an optimum temperature to survive.

The CEM recognizes canopy closure and local hydrology as habitat elements directly affecting temperature regulation. Disease may also have an effect.




Figure 7.—VEFL life stage 3 – breeding adult, basic CEM diagram. Only elements with connections within this life stage are presented.




Figure 8.—VEFL life stage 3 – breeding adult, high- and medium-magnitude relationships. Only elements with high- or medium-magnitude connections within this life stage are presented.

# Chapter 7 – Causal Relationships Across All Life Stages

The nine controlling factors discussed in chapter 5 have the same influence on the same habitat elements for all life stages for which those habitat elements matter. Table 5 shows the magnitudes of *direct* influence of the 9 controlling factors on the 15 habitat elements. The structure of table 5 is the same as for table 4, but table 5 shows the magnitudes of the relationships instead of just their presence/absence. The paragraphs following the table discuss the relative effects of the different controlling factors on each habitat element.

Table 5.—Magnitude of influence of controlling factors on habitat elements

Controlling factor →									
Habitat element affected ↓	Fire management	Grazing	Irrigation	Mechanical thinning	Nuisance species introduction and management	Pesticide/herbicide application	Planting regime	Recreational activities	Water storage-delivery system design and operation
Anthropogenic disturbance		Med.		High		Low	Low	Med.	
Brood size	N/A*								
Canopy closure	Med.			Med.	High		Med.	Med.	
Community type	Med.	Med.			High		Med.	Med.	Med.
Food availability					High	Med.	Med.		
Genetic diversity and infectious agents	N/A*								
Local hydrology			Med.						High
Parental feeding behavior								Med.	
Parental nest attendance								Med.	
Patch size	Med.	High		Med.	High		Med.	Med.	
Predator and cowbird density					Med.		Med.	Med.	
Soil salinity			Med.						Med.
Tree size	Med.			Low	Med.		Med.		
Understory density	Med.	Med.		Low	Med.		Low	Med.	Med.
Water availability			High						High

\* N/A values suggest that none of the identified controlling factors *directly* affect the habitat element.

## ANTHROPOGENIC DISTURBANCE

All activities involving humans increase anthropogenic disturbance. The scale and scope of the influences depend upon the scale and scope of the activity. In general, most activities are of narrow scope and short duration; however, systematic influences can cause repeated noise (e.g., campsites, off-highway vehicle trails, or nearby roads).

## CANOPY CLOSURE

The controlling factors that directly affect canopy closure include fire management, mechanical thinning, nuisance species introduction and management, planting regime, and recreational activities. Recreational activities, fire, and thinning will generally reduce canopy closure, whereas the effects of planting regime and nuisance species introduction and management depend on the management actions and species involved.

Fire affects many aspects of vegetation structure and composition and can destroy VEFL habitat (Engstrom et al. 1984). Fire management can have great effects on vegetation structure and is usually implemented over large areas. However, the dynamic nature of both fire and riparian communities means that effects of fire management will likely be short term in nature.

Mechanical thinning would be done at the patch level, with effects lasting until vegetation grows back, and can be as intense as managers wish.

Nuisance species can change the structure of entire communities, with lasting effects. Although effects are experienced at a patch level, invasive species can spread across entire regions, and their effects can last decades unless a complete transformation of the community type occurs.

Planting regimes have the ability to greatly affect vegetation. However, planting decisions are made at the scale of an individual restoration site. Although riparian communities tend to be ephemeral, restoration sites are heavily managed, so the effects are likely medium or even long term.

Finally, the potential impact of recreational activities on VEFL habitat is great, although it depends on the activity. Decisions regarding management of recreational activities can affect large areas, but the dynamic nature of both human activity and riparian communities means that effects of recreation will likely be short term in nature after the completion of the activity.

## COMMUNITY TYPE

The controlling factors that directly affect community type are fire management, grazing, nuisance species introduction and management, planting regime, recreational activities, and water storage-delivery system design and operation. It is not possible to state whether the effects of controlling factors are positive or negative on community type.

Fire affects many aspects of vegetation structure and composition and can destroy VEFL habitat (Engstrom et al. 1984). Fire management can have great effects on vegetation structure, and thus community type, and is usually implemented over large areas. However, the dynamic nature of both fire and riparian communities means that effects of fire management will likely be short term in nature.

Grazing affects many aspects of riparian vegetation structure and composition (Kauffman et al. 1997). Grazing activity can have great effects on community composition and is often implemented over large and long scales (Kauffman et al. 1997). However, the dynamic nature of riparian communities means that effects of grazing will likely be short term unless a complete transformation of the community type occurs. Krueper et al. (2003) note that VEFL recover quickly after the removal of cattle from the grazing system, indicating both the effect of grazing on VEFL presence and on the potential for recovery.

Nuisance species can completely change the structure of entire communities with permanent effects.

Planting regimes have the ability to greatly affect vegetation. However, planting decisions are made at the scale of individual restoration sites. Although riparian communities tend to be ephemeral, restoration sites are heavily managed, so the effects are likely medium or even long term.

Recreational activities can influence the species composition of riparian forests, although the effect depends on the activity.

Water storage and flow regimes can change the structure of entire plant communities with lasting effects (Nilsson and Svedmark 2002).

## FOOD AVAILABILITY

The controlling factors that directly affect the food available to VEFL include nuisance species introduction and management, pesticide/herbicide application, and planting regime.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

Nuisance species can change the arthropod community, as can the planting regime. The effects of nuisance species can spread across entire regions and result in a permanent transformation of the landscape.

The magnitude of the effect of pesticides/herbicides depends on many factors, but the potential magnitude is very high. However, the most likely scenario involves pesticide/herbicide applications at individual agricultural fields affecting nearby patches and the effects dissipating shortly after application. The diet of VEFL has not been studied but likely includes bees, wasps, flies, beetles, and grasshoppers (Rosenberg et al. 1991). VEFL have been observed feeding on fish on a few occasions, but this does not likely represent a major dietary item (Andrews et al. 1996).

## **LOCAL HYDROLOGY**

The only controlling factor affecting local hydrology is water storage-delivery system design and operation—it is not possible to put a direction on the effect. The amount of water released or stored affects water levels and therefore distance to water, soil moisture, and other hydrological conditions. Water storage and flow regimes can affect vegetation communities and food abundance (Nilsson and Svedmark 2002). The effects of water storage spreads over large scales, but the effects of changes in flow regimes likely will be short term in nature unless a complete transformation of the habitat occurs.

## **PARENTAL FEEDING BEHAVIOR**

Parental feeding behavior is influenced by any factors involving humans entering the nesting areas during the nesting phase. We have emphasized recreational activities as the most likely sustained disturbance that could be managed. Invasive species could play a role in disturbance if they included novel predators or parasites, but we chose not to include them, as none have been identified.

## **PARENTAL NEST ATTENDANCE**

Parental nest attendance is influenced by any factors involving humans entering the nesting areas during the nesting phase. We have emphasized recreational activities as the most likely sustained disturbance that could be managed. Invasive species could play a role in disturbance if they included novel predators or parasites, but we chose not to include them, as none have been identified.

## PATCH SIZE

The controlling factors that directly affect patch size include fire management, grazing, mechanical thinning, nuisance species introduction and management, planting regime, and recreational activities. Recreational activities, fire, grazing, and mechanical thinning will generally reduce the size of a given patch, whereas the effects of planting regime depend on the management actions and species involved.

Fire affects many aspects of vegetation structure composition and patch size and can destroy VEFL habitat (Engstrom et al. 1984). Fire management can have great effects on vegetation structure, and thus community type, and is usually implemented over large areas. However, the dynamic nature of both fire and riparian communities means that the effects of fire management will likely be short term in nature.

Grazing affects many aspects of riparian vegetation structure and composition (Kauffman et al. 1997). Grazing activity can have great effects on community composition, and is often implemented over large and long scales (Kauffman et al. 1997). However, the dynamic nature of riparian communities means that effects of grazing can be short-term in nature. Krueper et al. (2003) note that VEFL recover quickly after the removal of cattle from the grazing system, indicating both the effect of grazing on VEFL presence and on the potential for recovery.

Mechanical thinning is generally performed at the patch level, with effects lasting until vegetation grows back, and can be as intense as managers wish.

Nuisance species can change the structure of entire communities, with lasting effects. Although effects are experienced at a patch level, invasive species can spread across entire regions, and their effects can last decades unless a complete transformation of the community type occurs.

Planting regimes have the ability to greatly affect vegetation. However, planting decisions are made at the scale of individual restoration sites. Although riparian communities tend to be ephemeral, restoration sites are heavily managed, so the effects are likely medium or even long term.

Recreational activities can influence the species composition of riparian forests, although the effect depends on the activity. However, the dynamic nature of both human activity and riparian communities means that effects of recreation will likely be short term in nature after the completion of the activity.

## **PREDATOR AND COWBIRD DENSITY**

The controlling factors directly affecting predator and cowbird density include nuisance species introduction and management, planting regime, and recreational activities. The direction and size of these effects are difficult to quantify. Some studies have shown predator and cowbird presence differs among community types, native, and non-native habitats (Schmidt et al. 2005). Any change in the composition of the predator community can have a large and lasting impact on the VEFL population (Lima et al. 2009).

## **SOIL SALINITY**

The controlling factor directly affecting soil salinity is water storage-delivery system design and operation. Soil salinity is affected by the amount of water reaching the soil and the salinity of the water (San Joaquin River Restoration Program 2014). Main stem water generally has lower salinity levels than groundwater and thus can have a large impact on lowering soil salinity (San Joaquin River Restoration Program 2014).

## **TREE SIZE**

The controlling factors directly affecting tree size include fire management, mechanical thinning, nuisance species introduction and management, and planting regime. Fire and thinning will generally reduce tree size, whereas the effects of nuisance species introduction and management and planting regime depend on the management actions and species involved.

Fire affects many aspects of vegetation structure and composition and can destroy VEFL habitat (Engstrom et al. 1984). Fire management can have great effects on vegetation structure and can be implemented over small or large areas.

Mechanical thinning is generally performed at the patch level, with effects lasting until vegetation grows back, and can be as intense as managers wish.

Nuisance species can change the structure of entire communities, with lasting effects. Although effects are experienced at a patch level, invasive species can spread across entire regions, and their effects can last decades unless a complete transformation of the community type occurs. The effect on tree size depends upon the species of the invader and the species being displaced.

Planting regimes have the ability to greatly affect vegetation. The timing and type of trees planted can impact tree cover in the long run. However, planting

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

decisions are made at the scale of an individual restoration site. Although riparian communities tend to be ephemeral, restoration sites are heavily managed, so the effects are likely medium or even long term.

## **UNDERSTORY DENSITY**

The controlling factors that directly affect understory density include fire management, grazing, mechanical thinning, nuisance species introduction and management, planting regime, recreational activities, and water storage-delivery system design and operation. Recreational activities, fire, and mechanical thinning will generally reduce understory density, whereas the effects of nuisance species introduction and management, planting regime, and water storage-delivery system design and operation depend on the management actions and species involved.

Fire affects many aspects of vegetation structure and composition and can destroy VEFL habitat (Engstrom et al. 1984). Fire management can have great effects on vegetation structure and is usually implemented over large areas. However, the dynamic nature of both fire and riparian communities means that effects of fire management will likely last less than a decade.

Grazing affects many aspects of riparian vegetation structure and composition (Kauffman et al. 1997). Grazing activity can have great effects on community composition and is often implemented over large and long scales (Kauffman et al. 1997). However, the dynamic nature of riparian communities means that the effects of grazing will likely last less than a decade.

Mechanical thinning would be done at the patch level, with effects lasting until vegetation grows back, and can be as intense as managers choose. The effect only lasts until the vegetation grows back.

Nuisance species can change the structure of entire communities, with lasting effects. Although, effects are experienced at a patch level, invasive species can spread across entire regions, and their effects can last decades.

Planting regimes have the ability to greatly affect vegetation. However, planting decisions are made at the scale of individual restoration sites. Although riparian communities tend to be ephemeral, restoration sites are heavily managed, so the effects are likely medium or even long term.

The potential impact of recreational activities on VEFL habitat is great, although the effect depends on the activity. Decisions regarding management of recreational activities can affect large areas. However, the dynamic nature of both human activity and riparian communities means that effects of recreation will

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**

likely be short term in nature after the completion of the activity. Many recreational activities are growing within the area, so these effects could be long term in nature, and some may cause a permanent transition within the ecosystem.

Finally, the amount of water released or stored affects water levels and therefore distance to water, soil moisture, and other hydrological conditions. Water storage and flow regimes, specifically floods, can affect understory density (Hunter 1987a; Lite et al. 2005). The effects of flood spread over large scales, but the effects of changes in flow regimes likely last less than a decade unless a complete transformation of the habitat occurs.

## **WATER AVAILABILITY**

The presence of surface water is primarily influenced by main stem water management and by irrigation practices.

## Chapter 8 – Discussion and Conclusions


This chapter summarizes the findings of this assessment in three ways by posing three questions: (1) which critical biological activities and processes most strongly affect the individual across all life stages, (2) which habitat elements, in terms of their abundance, distribution, and quality, most strongly affect the most influential activities and processes, and (3) which of these causal relationships appear to be the least understood in ways that could affect their management?

### **MOST INFLUENTIAL ACTIVITIES AND PROCESSES ACROSS ALL LIFE STAGES**

Figure 9 identifies the critical biological activities and processes that the assessment found most strongly directly affect the success of VEFL at each life stage (high or medium magnitude). The findings presented in this diagram may be summarized as follows:

- Disease, eating/foraging, and predation are the most important critical biological activities processes affecting survival of VEFL at all life stages. Depredation of nests can be high and has been shown to be the primary cause of nest failure among some species along the LCR – exceeding 80 percent (Powell and Steidl 2000). The effects act at the landscape scale. Other processes such as molt and temperature regulation are important, but the effects are more indirect and less certain.
- Only two processes strongly and directly affect reproduction—nest attendance and nest site selection. These two critical biological activities and processes are especially important because they also affect the survival of the nestlings. Note, in this instance reproduction refers to the laying of eggs; the survival of the nestlings is modeled separately in our process.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**



**Figure 9.—Most influential biological activities and processes affecting each life stage of VEFL.**  
 Only elements with high- or medium-magnitude connections are presented. The legend is provided on figure 2.

## POTENTIALLY PIVOTAL ALTERATIONS TO HABITAT ELEMENTS

Figure 10 identifies the habitat elements that this assessment indicates most strongly directly affect the critical biological activities and processes identified on figure 9 across all life stages (high or medium magnitude). The findings presented in this diagram may be summarized as follows:

- Eating/foraging and nest site selection are by far affected by the most habitat variables. Eating/foraging has been the most studied among the factors influencing this species, yet much uncertainty remains. Eating/foraging is probably the top determinant as to whether individual birds choose to nest in the area.


**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

- Predation and brood parasitism are only strongly affected by two habitat elements—patch size and predator and cowbird density. Patch size affects both predation rates and brood parasitism rates because of its effects on the proportion of stand area near the edge of the patch, a factor which has been shown to influence predation (Brittingham and Temple 1996; Chalfoun and Martin 2009; Theimer et al. 2011). Predator and cowbird density affects predation rates (Lima 2009).
- Nest attendance is only strongly affected by brood size and predator and cowbird density. Brood size affects the amount of time adult VEFL must spend foraging versus attending the nest. Predator and cowbird density can influence nest defense behavior and thus nest attendance.
- Food availability and brood size drive the ability of VEFL at each life stage to acquire needed energy and nutrients.
- Disease and temperature regulation are important physiological concerns that can be impacted strongly by habitat elements such as canopy closure and local hydrology as well as the presence of genetic diversity and infectious agents.

## **GAPS IN UNDERSTANDING**

Figures 9 and 10 use the conventional color coding of individual causal relationships to identify relationships that the CEM identifies as having high, intermediate, or low levels of scientific confirmation. As noted in attachment 1, “Low” scientific understanding of a relationship means that it is “...subject to wide disagreement or uncertainty in peer-reviewed studies from within the ecosystem of concern and in scientific reasoning among experts familiar with the ecosystem.” In many cases, the scientific principles are well understood, but the factual details are insufficiently understood within the LCR. The two figures show large numbers of red arrows, indicating relationships that the assessment identifies as having a low level of scientific understanding. Each of these red arrows identifies a causal relationship that may warrant further field, laboratory, or literature investigation. The following paragraphs highlight some potentially important areas of low understanding.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model**



**Figure 10.—Habitat elements that directly affect the most influential biological activities and processes across all life stages of VEFL.** Only elements with high- or medium-magnitude connections within this life stage are presented. The legend is provided on figure 2.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

- Nest site selection is affected by the most habitat variables and is one of the most well-understood processes. However, many gaps in knowledge exist, and the sensitivity of the species to subtle changes in the nest stands is not thoroughly understood.
- The effects of predation on juveniles and adults is poorly understood, whereas nest predation is better studied. This likely reflects the relative ease of studying depredation of nests versus free-flying birds. If the persistence or population growth of VEFL populations is considered sensitive to the survival of adults and juveniles, then research regarding predation should be a considered.
- The indirect influences on food availability have a large impact on eating/foraging, which is then directly linked with survival of VEFL at all life stages. Further investigation into these indirect effects is warranted.
- Anthropogenic disturbance has been noted to have a broad range of impacts on the ecology of birds (Francis and Barber 2013). Noise has been shown to affect foraging efficiency in many species but generally affects different species in different ways. For example, VEFL have been observed to sing longer songs in noisier environments (Ríos-Chelén et al. 2012). The effects of noise within the LCR have not been studied and are difficult to quantify.
- The effects of disease, ecto-parasites, and endo-parasites have not been studied in the VEFL or among passerine species inhabiting the LCR. Diseases have the potential to have dramatic impacts on populations (Robinson et al. 2010). Further, the genetic health of VEFL within the LCR has not been evaluated.

This list of uncertainties is not meant to be exhaustive but only to highlight topics the literature identifies as potentially pivotal to VEFL recruitment along the LCR and to identify important gaps in these publications. They are not in any way to be considered guidance for Reclamation or LCR MSCP, nor are these knowledge gaps expected to be addressed under the program.

## LITERATURE CITED

- Andrews, B.J., M. Sullivan, and J.D. Hoerath. 1996. Vermilion flycatcher and black phoebe feeding on fish. *The Wilson Bulletin* 108:377–378.
- Archer, T.J. 1996. Observations on nesting and display flights of the vermilion flycatcher in western Texas. *The Southwestern Naturalist* 41:443–444.
- Barber, J.R., K.R. Crooks, and K.M. Fristrup. 2010. The costs of chronic noise exposure for terrestrial organisms. *Trends in Ecology & Evolution* 25:180–189.
- Bateman, H.L., P.L. Nagler, and E.P. Glenn. 2013. Plot- and landscape-level changes in climate and vegetation following defoliation of exotic saltcedar (*Tamarix* sp.) from the biocontrol agent *Diorhabda carinulata* along a stream in the Mojave Desert (USA). *Journal of Arid Environments* 89:16–20.
- Birkhead, T.R., F. Fletcher, and E.J. Pellatt. 1999. Nestling diet, secondary sexual traits and fitness in the Zebra Finch. *Proceedings: Biological Sciences* 266:385–390.
- Boyle, S.A. and F.B. Samson. 1985. Effects of nonconsumptive recreation on wildlife: a review. *Wildlife Society Bulletin* 13:110–116.
- Brand, L.A., J.C. Stromberg, and B.R. Noon. 2010. Avian density and nest survival on the San Pedro River: importance of vegetation type and hydrologic regime. *The Journal of Wildlife Management* 74:739–754.
- Brittingham, M.C. and S.A. Temple. 1996. Vegetation around parasitized and non-parasitized nests within deciduous forest. *Journal of Field Ornithology* 67:406–413.
- Burke, M., K. Jorde, and J.M. Buffington. 2009. Application of a hierarchical framework for assessing environmental impacts of dam operation: changes in streamflow, bed mobility and recruitment of riparian trees in a western North American river. *Journal of Environmental Management* 90:S224–S236.
- Carothers, S.W. 1974. Breeding ecology and time-energy budget of male vermilion flycatchers and comments on the social organization of southwestern riparian birds. Ph.D. thesis. University of Illinois at Urbana-Champaign.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**

- Chalfoun, A.D. and T.E. Martin. 2009. Habitat structure mediates predation risk for sedentary prey: experimental tests of alternative hypotheses. *The Journal of Animal Ecology* 78:497–503.
- Di Tomaso, J.M. 1998. Impact, biology, and ecology of saltcedar (*Tamarix* spp.) in the southwestern United States. *Weed technology* 12:326–336.
- DiGennaro, B., D. Reed, C. Swanson, L. Hastings, Z. Hymanson, M. Healey, S. Siegel, S. Cantrell, and B. Herbold. 2012. Using conceptual models and decision-support tools to guide ecosystem restoration planning and adaptive management: an example from the Sacramento–San Joaquin Delta, California. *San Francisco Estuary and Watershed Science* 10:1–15.
- Ellison, K.S. 2008. Nest reuse by vermilion flycatchers in Texas. *Wilson Journal of Ornithology* 120:339–344.
- Ellison, K.S., S.G. Sealy, and H.R. McGaha. 2007. Color variation among nestling brown-headed cowbirds (*Molothrus ater*) does not reflect differential success with hosts in Texas. *Auk* 124:526–536
- Ellison, K.S., B.O. Wolf, and S.L. Jones. 2009. Vermilion Flycatcher (*Pyrocephalus rubinus*). *The Birds of North America Online*. Ithaca, New York: Cornell Lab of Ornithology.  
<http://bna.birds.cornell.edu/bna/species/484>
- Engstrom, R.T., R.L. Crawford, and W.W. Baker. 1984. Breeding bird populations in relation to changing forest structure following fire exclusion: a 15-year study. *Wilson Bulletin* 96:437–450.
- Etterson, M.A., S.N. Ellis-Felege, D. Evers, G. Gauthier, J.A. Grzybowski, B.J. Mattsson, L.R. Nagy, B.J. Olsen, C.M. Pease, M.P. van der Burg, and A. Potvien. 2011. Modeling fecundity in birds: conceptual overview, current models, and considerations for future developments. *Ecological Modelling* 222:2178–2190.
- Fischenich, J.C. 2008. The application of conceptual models to ecosystem restoration. Vicksburg, Mississippi: U.S. Army Engineer Research and Development Center. Technical Note EMRRP-EBA-01.
- Francis, C.D. and J.R. Barber. 2013. A framework for understanding noise impacts on wildlife: an urgent conservation priority. *Frontiers in Ecology and the Environment* 11:305–313.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

- Great Basin Bird Observatory. 2010. Summary Report on the Lower Colorado River Riparian Bird Surveys, 2008–2010. Unpublished report submitted to the L Bureau of Reclamation, Lower Colorado River Multi-Species Conservation Program, Boulder City, Nevada, by the Great Basin Bird Observatory, Reno, Nevada.
- Howell, S.N.G. 2010. Molt in North American Birds. Boston, Massachusetts: Houghton Mifflin Harcourt.
- Hunter, W.C., B.W. Anderson, and R.D. Ohmart. 1987a. Avian community structure changes in a mature floodplain forest after extensive flooding. *The Journal of Wildlife Management* 51:495–502.
- Hunter, W.C., R.D. Ohmart, and B.W. Anderson. 1987b. Status of breeding riparian-obligate birds in southwestern riverine systems. *Western Birds* 18:10–18.
- Jarvi, S.I., J.J. Schultz, and C.T. Atkinson. 2002. PCR diagnostics underestimate the prevalence of avian malaria (*Plasmodium relictum*) in experimentally-infected passerines. *The Journal of Parasitology* 88:153–159.
- Jennings, S.B., N.D. Brown, and D. Sheil. 1999. Assessing forest canopies and understorey illumination: canopy closure, canopy cover and other measures. *Forestry* 72:59–74.
- Kauffman, J.B., R.L. Beschta, N. Otting, and D. Lytjen. 1997. An ecological perspective of riparian and stream restoration in the western United States. *Fisheries* 22:12–24.
- Klimkiewicz, M. K., and A.G. Futcher. 1989. Longevity records of North American birds, supplement 1. *Journal of Field Ornithology* 60:469–494.
- Kondolf, G.M., J.G. Williams, T.C. Horner, and D. Milan. 2008. Assessing physical quality of spawning habitat. Pages 249–274 in D.A Sear and P. DeVries (editors). *Salmonid Spawning Habitat in Rivers: Physical Controls, Biological Responses, and Approaches*. American Fisheries Society Symposium 65. American Fisheries Society, Bethesda, Maryland.
- Krueper D., J. Bart, and T.D. Rich. 2003. Response of vegetation and breeding birds to the removal of cattle on the San Pedro River, Arizona (U.S.A.). *Conservation Biology* 17:607–615.
- Lachish, S., S.C.L. Knowles, R. Alves, M.J. Wood, and B.C. Sheldon. 2011. Fitness effects of endemic malaria infections in a wild bird population: the importance of ecological structure. *The Journal of Animal Ecology* 80:1196–1206.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)**  
**Basic Conceptual Ecological Model for the Lower Colorado River**

- Landres, P.B. and J.A. MacMahon. 1980. Guilds and community organization: analysis of an oak woodland avifauna in Sonora, Mexico. *Auk* 97:351–365.
- Lima, S.L. 1998. Nonlethal effects in the ecology of predator-prey interactions. *BioScience* 48:25–34.
- \_\_\_\_\_. 2009. Predators and the breeding bird: behavioral and reproductive flexibility under the risk of predation. *Biological Reviews of the Cambridge Philosophical Society* 84:485–513.
- Lite, S.J., K.J. Bagstad, and J.C. Stromberg. 2005. Riparian plant species richness along lateral and longitudinal gradients of water stress and flood disturbance, San Pedro River, Arizona, USA. *Journal of Arid Environments* 63:785–813.
- Mason, P. 1986. Brood parasitism in a host generalist, the Shiny Cowbird: I. The quality of different species as hosts. *The Auk* 103:52–60.
- McDonald, D.B. and H. Caswell. 1993. Matrix methods for avian demography. *Current Ornithology* 10:139–185.
- Merino, S., J. Moreno, J.J. Sanz, and E. Arriero. 2000. Are avian blood parasites pathogenic in the wild? A medication experiment in Blue Tits (*Parus caeruleus*). *Proceedings: Biological Sciences* 267:2507–2510.
- Michel, N., D.F. DeSante, D.R. Kaschube, and M.P. Nott. 2006. The Monitoring Avian Productivity and Survivorship (MAPS) Program Annual Reports, 1989–2003. The Institute for Bird Populations, Point Reyes Station, California.
- Morishita, T.Y., P.P. Aye, E.C. Ley, and B.S. Harr. 1999. Survey of pathogens and blood parasites in free-living passerines. *Avian Diseases* 43:549–552.
- Myers, S.J. 2008. Vermilion flycatcher (*Pyrocephalus rubinus*) in W.D. Shuford and T. Gardali (editors). *California Bird Species of Special Concern: A Ranked Assessment of Species, Subspecies, and Distinct Populations of Birds of Immediate Conservation Concern in California*. Western Field Ornithologists, Camarillo, California, and California Department of Fish and Game, Sacramento.
- Nilsson, C. and M. Svedmark. 2002. Basic principles and ecological consequences of changing water regimes: riparian plant communities. *Environmental Management* 30:468–480.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

- Palinauskas, V., G. Valkiūnas, C.V. Bolshakov, and S. Bensch. 2008. *Plasmodium relictum* (lineage P-SGS1): effects on experimentally infected passerine birds. *Experimental Parasitology* 120:372–80.
- Phillips, A., J. Marshall, and G. Monson. 1964. *The Birds of Arizona*. Tucson, Arizona: University of Arizona Press.
- Powell, B.F. and R.J. Steidl. 2000. Nesting habitat and reproductive success of Southwestern riparian birds. *The Condor* 102:823–831.
- \_\_\_\_\_. 2002. Habitat selection by riparian songbirds breeding in southern Arizona. *The Journal of Wildlife Management* 66:1096–1103.
- Rasmussen, J.L. and S.G. Sealy. 2006. Hosts feeding only Brown-headed Cowbird fledglings: where are the host fledglings? *Journal of Field Ornithology* 77:269–279.
- Raulston, B. 2013. Vermilion Flycatchers on the LCR: A Summary of Data From 1970–2012. Lower Colorado River Multi-Species Conservation Program, Bureau of Reclamation, Boulder City, Nevada.
- Ricklefs, R.E. and R.E. Shea. 2007. Estimating annual survival in sexually dimorphic species from proportions of first-year birds. *Ecology* 88:1408–1419.
- Ríos-Chelén, A.A., J.A. Graves, R. Torres, M. Serrano-Pinto, L. D’Alba, and C. M. García. 2008. Intra-specific brood parasitism revealed by DNA micro-satellite analyses in a sub-oscine bird, the vermilion flycatcher. *Revista Chilena de Historia Natural* 81:21–31.
- Ríos-Chelén, A.A., E. Quirós-Guerrero, D. Gil, and C. Macías Garcia. 2012. Dealing with urban noise: Vermilion Flycatchers sing longer songs in noisier territories. *Behavioral Ecology and Sociobiology* 67:145–152.
- Robinson, R.A., B. Lawson, M.P. Toms, K.M. Peck, J.K. Kirkwood, J. Chantrey, I.R. Clatworthy, A.D. Evans, L.A. Hughes, O.C. Hutchinson, S.K. John, T.W. Pennycott, M.W. Perkins, P.S. Rowley, V.R. Simpson, K.M. Tyler, and A.A. Cunningham. 2010. Emerging infectious disease leads to rapid population declines of common British birds. *PloS One* 5:e12215.
- Rosenberg, K.V., R.D. Ohmart, W.C. Hunter, and B.W. Anderson. 1991. *Birds of the Lower Colorado River Valley*. Tucson: University of Arizona Press.
- Rosenberg, K.V., J.D. Lowe, and A.A. Dhondt. 1999. Effects of forest fragmentation on breeding tanagers: a continental perspective. *Conservation Biology* 13:568–583.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

- Rosenfeld, J.S. 2003. Assessing the habitat requirements of stream fishes: an overview and evaluation of different approaches. *Transactions of the American Fisheries Society* 132:953–968.
- Rosenfeld, J.S. and T. Hatfield. 2006. Information needs for assessing critical habitat of freshwater fish. *Canadian Journal of Fisheries and Aquatic Sciences* 63:683–698.
- San Joaquin River Restoration Program. 2014. Seepage Management Plan (Draft September 2014).  
<http://www.restoresjr.net/>
- Schmidt, K.A., L.C. Nelis, N. Briggs, and R.S. Ostfeld. 2005. Invasive shrubs and songbird nesting success: effects of climate variability and predator abundance. *Ecological Applications* 15:258–265.
- Small, A. 1994. *California Birds: Their Status and Distribution*. Ibis Publishing Company, Vista, California. 342 p.
- Taylor, W.K. and H. Hanson. 1970. Observations on the breeding biology of the vermilion flycatcher in Arizona. *The Wilson Bulletin* 82:315–319.
- Theimer, T., D. Peterson, A. Pellegrini, M.A. McLeod, and T. Koronkiewicz. 2011. Real and artificial nest predation and parental nest attendance along the Lower Colorado River and southern Nevada. U.S. Department of the Interior, Bureau of Reclamation, Lower Colorado Regional Office, Boulder City, Nevada.
- U.S. Fish and Wildlife Service (USFWS). 2002. *Southwestern Willow Flycatcher Recovery Plan*, Albuquerque, New Mexico. U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- \_\_\_\_\_. 2013. Designation of critical habitat for southwestern willow flycatcher; final rule. *Federal Register* 78(2):344–534.
- Ware, H.E., C.J.W. McClure, J.D. Carlisle, and J.R. Barber. 2015. A phantom road experiment reveals traffic noise is an invisible source of habitat degradation. *Proceedings of the National Academy of Sciences*, 112:12105–12109.
- Wildhaber, M.L., A.J. DeLonay, D.M. Papoulias, D.L. Galat, R.B. Jacobson, D.G. Simpkins, P.J. Baaten, C.E. Korschgen, and M.J. Mac. 2007. A conceptual life-history model for pallid and shovelnose sturgeon. U.S. Geological Survey, Circular 1315. Reston, Virginia.

**Vermilion Flycatcher (*Pyrocephalus rubinus*) (VEFL)  
Basic Conceptual Ecological Model for the Lower Colorado River**

- \_\_\_\_\_. 2011. Identifying structural elements needed for development of a predictive life-history model for pallid and shovelnose sturgeons. *Journal of Applied Ichthyology* 27:462–469.
- Winfree, R. 2004. High offspring survival of the brown-headed cowbird in an invaded habitat. *Animal Conservation* 7:445–453.

## ACKNOWLEDGMENTS

The authors would like to acknowledge Barb Raulston, Beth Sabin, and Chris Dodge, biologists with Reclamation, LCR MSCP; and Sonja Kokos, Adaptive Management Group Manager, LCR MSCP, who provided invaluable technical feedback and guidance during the development of the model process and production of this report. We would also like to acknowledge John Swett, Program Manager, LCR MSCP, for his leadership and support of this modeling effort that will guide and inform the work of the LCR MSCP well into the future.

# **ATTACHMENT 1**

Species Conceptual Ecological Model Methodology for the  
Lower Colorado River Multi-Species Conservation Program

# OVERVIEW OF METHODOLOGY

The conceptual ecological models (CEMs) for species covered by the Lower Colorado River Multi-Species Conservation Program (LCR MSCP) Habitat Conservation Plan expand on a methodology developed by the Sacramento-San Joaquin Delta Ecosystem Restoration Program (ERP): [https://www.dfg.ca.gov/ERP/conceptual\\_models.asp](https://www.dfg.ca.gov/ERP/conceptual_models.asp). The ERP is jointly implemented by the California Department of Fish and Wildlife, U.S. Fish and Wildlife Service, and National Marine Fisheries Service. The Bureau of Reclamation participates in this program.

The ERP methodology incorporates common best practices for constructing CEMs for individual species (Wildhaber et al. 2007; Fischenich 2008; DiGennaro et al. 2012). It has the following key features:

- It focuses on the *major life stages or events* through which each species passes and the *output(s)* of each life stage or event. Outputs typically consist of survivorship or the production of offspring.
- It identifies the *major drivers* that affect the likelihood (rate) of each output. Drivers are physical, chemical, or biological factors – both natural and anthropogenic – that affect output rates and therefore control the viability of the species in a given ecosystem.
- It characterizes these interrelationships using a “*driver-linkage-outcomes*” approach. Outcomes are the output rates. Linkages are cause-effect relationships between drivers and outcomes.
- It *characterizes each causal linkage* along four dimensions: (1) the character and direction of the effect, (2) the magnitude of the effect, (3) the predictability (consistency) of the effect, and (4) the certainty of present scientific understanding of the effect (DiGennaro et al. 2012).

The CEM methodology used for species covered by the LCR MSCP Habitat Conservation Plan species expands this ERP methodology. Specifically, the present methodology incorporates the recommendations and examples of Wildhaber et al. (2007, 2011), Kondolf et al. (2008), and Burke et al. (2009) for a more hierarchical approach and adds explicit demographic notation for the characterization of life-stage outcomes (McDonald and Caswell 1993). This expanded approach provides greater detail on causal linkages and outcomes. The expansion specifically calls for identifying **four** types of model components for each life stage, and the causal linkages among them, as follows:

- **Life-stage outcomes** are outcomes of an individual life stage, including the recruitment of individuals to the next succeeding life stage (e.g., juvenile to adult). For some life stages, the outcomes, alternatively or additionally, may include the survival of individuals to an older age class within the same life stage or the production of offspring. The rates of life-stage outcomes depend on the rates of the critical biological activities and processes for that life stage.
- **Critical biological activities and processes** are activities in which a species engages and the biological processes that must take place during each life stage that significantly affect life-stage outcomes. They include activities and processes that may benefit or degrade life-stage outcomes. Examples of critical activities and processes include mating, foraging, avoiding predators, avoiding other specific hazards, gamete production, egg maturation, leaf production, and seed germination. Critical activities and processes are “rate” variables. Taken together, the rate (intensity) of these activities and processes determine the rates of different life-stage outcomes.
- **Habitat elements** are specific habitat conditions that significantly ensure, allow, or interfere with critical biological activities and processes. The full suite of natural habitat elements constitutes the natural habitat template for a given life stage. Human activities may introduce habitat elements not present in the natural habitat template. Defining a habitat element may involve estimating the specific ranges of quantifiable properties of that element *whenever the state of knowledge supports such estimates*. These properties concern the abundance, spatial and temporal distributions, and other qualities of the habitat element that significantly affect the ways in which it ensures, allows, or interferes with critical biological activities and processes.
- **Controlling factors** are environmental conditions and dynamics – both natural and anthropogenic – that determine the quality, abundance, and spatial and temporal distributions of one or more habitat elements. In some instances, a controlling factor alternatively or additionally may directly affect a critical biological activity or process. Controlling factors are also called “drivers.” A hierarchy of controlling factors will exist, affecting the system at different temporal and spatial scales. Long-term dynamics of climate and geology define the domain of this hierarchy (Burke et al. 2009). For example, the availability of suitable nest sites for a riparian nesting bird may depend on factors such as canopy closure, community type, humidity, and intermediate structure which, in turn, may depend on factors such as water storage-delivery system design and operation (dam design, reservoir morphology, and dam operations) which, in turn, is shaped by watershed geology, vegetation, climate, land use, and water demand. *The LCR MSCP conceptual ecological models focus*

*on controlling factors that are within the scope of potential human manipulation, including management actions directed toward the species of interest.*

The present CEM methodology also explicitly defines a “life stage” as a biologically distinct portion of the life cycle of a species. The individuals in each life stage undergo distinct developments in body form and function; engage in distinct types behaviors, including reproduction; use different sets of habitats or the same habitats in different ways; interact differently with their larger ecosystems; and/or experience different types and sources of stress. A single life stage may include multiple age classes. A CEM focused on life stages is not a demographic model per se (McDonald and Caswell 1993). Instead, it is a complementary model focused on the ecological factors (drivers) that shape population dynamics.

This expanded approach permits the consideration of **six** possible types of causal relationships, on which management actions may focus, for each life stage of a species:

- (1) The effect of one controlling factor on another
- (2) The effect of a controlling factor on the abundance, spatial and temporal distributions, and other qualities of a habitat element
- (3) The effect of the abundance, spatial and temporal distributions, and other qualities of one habitat element on those of another
- (4) The effect of the abundance, spatial and temporal distributions, and other qualities of a habitat element on a critical biological activity or process
- (5) The effect of one critical biological activity or process on another
- (6) The effect of a critical biological activity or process on a specific life-stage outcome

Each controlling factor may affect the abundance, spatial and temporal distributions, and other qualities of more than one habitat element and several controlling factors may affect the abundance, spatial or temporal distributions, or other qualities of each habitat element. Similarly, the abundance, spatial and temporal distributions, and other qualities of each habitat element may affect more than one biological activity or process, and the abundances, spatial or temporal distributions, or other qualities of several habitat elements may affect each biological activity or process. Finally, the rate of each critical biological activity or process may contribute to the rates of more than one life-stage outcome.

Integrating this information across all life stages for a species provides a detailed picture of: (1) what is known, with what certainty, and the sources of this information; (2) critical areas of uncertain or conflicting science that demand resolution to better guide LCR MSCP management planning and action; (3) crucial attributes to use to monitor system conditions and predict the effects of experiments, management actions, and other potential agents of change; and (4) how managers may expect the characteristics of a resource to change as a result of changes to controlling factors, including changes in management actions.

## **Conceptual Ecological Models as Hypotheses**

The CEM for each species produced with this methodology constitutes a collection of hypotheses for that species. These hypotheses concern: (1) the species' life history; (2) the species' habitat requirements and constraints; (3) the factors that control the quality, abundance, and spatial and temporal distributions of these habitat conditions; and (4) the causal relationships among these. Knowledge about these model components and relationships may vary, ranging from well settled to very tentative. Such variation in the certainty of current knowledge always arises as a consequence of variation in the types and amount of evidence available and in the ecological assumptions applied by different experts.

Wherever possible, the information assembled for the LCR MSCP species CEMs documents the degree of certainty of current knowledge concerning each component and linkage in the model. This certainty is indicated by the quality, abundance, and consistency of the available evidence and by the degree of agreement/disagreement among the experts. Differences in the interpretations or arguments offered by different experts may be represented as alternative hypotheses. Categorizing the degree of agreement/disagreement concerning the components and linkages in a CEM makes it easier to identify topics of greater uncertainty or controversy.

## **Characterizing Causal Relationships**

A causal relationship exists when a change in one condition or property of a system results in a change in some other condition or property. A change in the first condition is said to cause a change in the second condition. The present CEM methodology includes methods for assessing causal relationships (links) along four dimensions (attributes) adapted from the ERP methodology (DiGennaro et al. 2012):

- (1) The character and direction of the effect
- (2) The magnitude of the effect
- (3) The predictability (consistency) of the effect
- (4) The certainty of present scientific understanding of the effect

The present and ERP methodologies for assessing causal linkages differ in three ways. First, the ERP methodology assesses these four attributes for the *cumulative* effect of the entire causal chain leading up to each outcome. However, the LCR MSCP methodology recognizes six different types of causal linkages as described above. This added level of detail and complexity makes it difficult in a single step to assess the cumulative effects of all causal relationships that lead up to any one individual causal link. For example, in the present methodology, the effect of a given critical biological activity or process on a particular life-stage outcome may depend on the effects of several habitat elements on that critical biological activity or process which, in turn, may depend on the effects of several controlling factors. For this reason, the present methodology assesses the four attributes separately for each causal link *by itself* rather than attempting to assess cumulative effects of all causal linkages leading to the linkage of interest. The present methodology assesses cumulative effects instead through analyses of the data assembled on all individual linkages. The analyses are made possible by assembling the data on all individual linkages in a spreadsheet as described below.

Second, the present CEM methodology explicitly divides link magnitude into three separate subattributes and provides a specific methodology for integrating their rankings into an overall ranking for link magnitude: (1) link intensity, (2) link spatial scale, and (3) link temporal scale. In contrast, the ERP methodology treats spatial and temporal scale together and does not separately evaluate link intensity. The present methodology defines link intensity as the relative strength of the effect of the causal node on the affected node *at the places and times where the effect occurs*. Link spatial scale is the relative spatial extent of the effect of the causal node on the affected node. Link temporal scale is the relative temporal extent of the effect of the causal node on the affected node. The present methodology defines link magnitude as the average of the separate rankings of link intensity, spatial scale, and temporal scale as described below.

Third, the ERP methodology addresses a single, large landscape, while the present methodology needed the flexibility to generate models applicable to a variety of spatial scopes. For example, the present methodology needed to support modeling of a single restoration site, the LCR main stem and flood plain, or the entire Lower Colorado River Basin. Consequently, the present methodology assesses the spatial scale of cause-effect relationships only relative to the spatial scope of the model.

The LCR MSCP conceptual ecological model methodology thus defines the four attributes for a causal link as follows:

- **Link character** – This attribute categorizes a causal relationship as positive, negative, involving a threshold response, or “complex.” “Positive” means that an increase in the causal node results in an increase in the affected node, while a decrease in the causal node results in a decrease in the affected node. “Negative” means that an increase in the causal node results in a decrease in the affected element, while a decrease in the causal node results in an increase in the affected node. Thus, “positive” or “negative” here do *not* mean that a relationship is beneficial or detrimental. The terms instead provide information analogous to the sign of a correlation coefficient. “Threshold” means that a change in the causal agent must cross some value before producing an effect. “Complex” means that there is more going on than a simple positive, negative, or threshold effect. In addition, this attribute categorizes a causal relationship as uni- or bi-directional. Bi-directional relationships involve a reciprocal relationship in which each node affects the other.
- **Link magnitude** – This attribute refers to “... the degree to which a linkage controls the outcome *relative to other drivers*” (DiGennaro et al. 2012). Magnitude takes into account the spatial and temporal scale of the causal relationship as well as the strength (intensity) of the relationship in individual locations. The present methodology provides separate ratings for the intensity, spatial scale, and temporal scale of each link, as defined above, and assesses overall link magnitude by averaging these three elements. Just as the terms for link character provide information analogous to the sign of a correlation coefficient, the terms for link magnitude provide information analogous to the size of a correlation coefficient. Tables 1-1 through 1-4 present the rating framework for link magnitude.
- **Link predictability** – This attribute refers to “... the degree to which the current understanding of the system can be used to predict the role of the driver in influencing the outcome. Predictability ... captures variability ... [and recognizes that] effects may vary so much that properly measuring and statistically characterizing inputs to the model are difficult” (DiGennaro et al. 2012). A causal relationship may be unpredictable because of natural variability in the system or because its effects depend on the interaction of other factors with independent sources for their own variability. Just as the terms for link character provide information analogous to the sign of a correlation coefficient, the terms for link predictability provide information analogous to the size of the range of error for a correlation coefficient. Table 1-5 presents the scoring framework for link predictability.

- **Link understanding** refers to the degree of agreement represented in the scientific literature and among experts in understanding how each driver is linked to each outcome. Table 1-6 presents the scoring framework for understanding. Link predictability and understanding are independent attributes. A link may be considered highly predictable but poorly understood or poorly predictable but well understood.

## Conceptual Ecological Model Documentation

The documentation for each CEM provides information in three forms: (1) a narrative report, (2) causal diagrams showing the model components and their causal linkages for each life stage, and (3) a spreadsheet that is used to record the detailed information (e.g., linkage attribute ratings) for each causal linkage. The spreadsheet and diagrams, built using Microsoft Excel™ and Microsoft Visio™, respectively, are linked so that the diagrams provide a fully synchronized summary of the information in the spreadsheet.

The narrative report for each species presents the definitions and rationales for the life stages/events and their outcomes identified for the species' life history; the critical biological activities and processes identified for each life stage; the habitat elements identified as supporting or impeding each critical biological activity or process for each life stage; the controlling factors identified as affecting the abundance, spatial and temporal distributions, and other qualities of the habitat elements for each life stage; and the causal linkages among these model components.

The narrative report includes causal diagrams (*aka* “influence diagrams”) for each life stage. These diagrams show the individual components or nodes of the model for that stage (life-stage outcomes, critical biological activities and processes, habitat elements, and controlling factors) and their causal relationships. The causal relationships (causal links) are represented by arrows indicating which nodes are linked and the directions of the causal relationships. The attributes of each causal link are represented by varying line thickness, line color, and other visual properties as shown on figure 1-1. The diagram conventions mostly follow those in the ERP methodology (DiGennaro et al. 2012).

The spreadsheet for each CEM contains a separate worksheet for each life stage. Each row in the worksheet for a life stage represents a single causal link. Table 1-7 lists the fields (columns) recorded for each causal link.

## Link Attribute Ratings, Spreadsheet Fields, and Diagram Conventions

Table 1-1.—Criteria for rating the relative intensity of a causal relationship – one of three variables in the rating of link magnitude (after DiGennaro et al. 2012, Table 2)

<b>Link intensity</b> – the relative strength of the effect of the causal node on the affected node <i>at the places and times where the effect occurs.</i>	
High	Even a relatively small change in the causal node will result in a relatively large change in the affected node <i>at the places and times where the effect occurs.</i>
Medium	A relatively large change in the causal node will result in a relatively large change in the affected node; a relatively moderate change in the causal node will result in no more than a relatively moderate change in the affected node; and a relatively small change in the causal node will result in no more than a relatively small change in the affected node <i>at the places and times where the effect occurs.</i>
Low	Even a relatively large change in the causal node will result in only a relatively small change in the affected node <i>at the places and times where the effect occurs.</i>
Unknown	Insufficient information exists to rate link intensity.

Table 1-2.—Criteria for rating the relative spatial scale of a cause-effect relationship – one of three variables in the rating of link magnitude (after DiGennaro et al. 2012, Table 1)

<b>Link spatial scale</b> – the relative spatial extent of the effect of the causal node on the affected node. The rating takes into account the spatial scale of the cause and its effect.	
Large	Even a relatively small change in the causal node will result in a change in the affected node across a large fraction of the spatial scope of the model.
Medium	A relatively large change in the causal node will result in a change in the affected node across a large fraction of the spatial scope of the model; a relatively moderate change in the causal node will result in a change in the affected node across no more than a moderate fraction of the spatial scope of the model; and a relatively small change in the causal node will result in a change in the affected node across no more than a small fraction of the spatial scope of the model.
Small	Even a relatively large change in the causal node will result in a change in the affected node across only a small fraction of the spatial scope of the model.
Unknown	Insufficient information exists to rate link spatial scale.

Table 1-3.—Criteria for rating the relative temporal scale of a cause-effect relationship – one of three variables in the rating of link magnitude (after DiGennaro et al. 2012, Table 1)

<b>Link temporal scale</b> – the relative temporal extent of the effect of the causal node on the affected node. The rating takes into account the temporal scale of the cause and its effect.	
Large	Even a relatively small change in the causal node will result in a change in the affected node that persists or recurs over a relatively large span of time – decades or longer – even without specific intervention to sustain the effect.
Medium	A relatively large change in the causal node will result in a change in the affected node that persists or recurs over a relatively large span of time – decades or longer – even without specific intervention to sustain the effect; a relatively moderate change in the causal node will result in a change in the affected node that persists or recurs over only a relatively moderate span of time – one or two decades – without specific intervention to sustain the effect; a relatively small change in the causal node will result in a change in the affected node that persists or recurs over only a relatively short span of time – less than a decade – without specific intervention to sustain the effect.
Small	Even a relatively large change in the causal node will result in a change in the affected node that persists or recurs over only a relatively short span of time – less than a decade – without specific intervention to sustain the effect.
Unknown	Insufficient information exists to rate link temporal scale.

Table 1-4.—Criteria for rating the overall relative link magnitude of a cause-effect relationship based on link intensity, spatial scale, and temporal scale

<b>Link magnitude</b> – the overall relative magnitude of the effect of the causal node on the affected node based on the numerical average for link intensity, spatial scale, and temporal scale. (Calculated by assigning a numerical value of 3 to “High” or “Large,” 2 to “Medium,” 1 to “Low” or “Small,” and not counting missing or “Unknown” ratings.)	
High	Numerical average $\geq 2.67$
Medium	Numerical average $\geq 1.67$ but $< 2.67$
Low	Numerical average $< 1.67$
Unknown	No subattribute is rated High/Large, Medium, or Low/Small, but at least one subattribute is rated Unknown.

Table 1-5.—Criteria for rating the relative predictability of a cause-effect relationship (after DiGennaro et al. 2012, Table 3)

<b>Link predictability</b> – the statistical likelihood that a given causal agent will produce the effect of interest.	
High	Magnitude of effect is largely unaffected by random variation or by variability in other ecosystem dynamics or external factors.
Medium	Magnitude of effect is moderately affected by random variation or by variability in other ecosystem processes or external factors.
Low	Magnitude of effect is strongly affected by random variation or by variability in other ecosystem processes or external factors.
Unknown	Insufficient information exists to rate link predictability.

Table 1-6.—Criteria for rating the relative understanding of a cause-effect relationship (after DiGennaro et al. 2012, Table 3)

<b>Understanding</b> – the degree of agreement in the literature and among experts on the magnitude and predictability of the cause-effect relationship of interest.	
High	Understanding of the relationship is subject to little or no disagreement or uncertainty in peer-reviewed studies from within the ecosystem of concern or in scientific reasoning among experts familiar with the ecosystem. Understanding may also rest on well-accepted scientific principles and/or studies in highly analogous systems.
Medium	Understanding of the relationship is subject to moderate disagreement or uncertainty in peer-reviewed studies from within the ecosystem of concern and in scientific reasoning among experts familiar with the ecosystem.
Low	Understanding of the relationship is subject to wide disagreement, uncertainty, or lack of evidence in peer-reviewed studies from within the ecosystem of concern and in scientific reasoning among experts familiar with the ecosystem.
Unknown	<i>(The “Low” rank includes this condition).</i>

Table 1-7.—Organization of the worksheet for each life stage

Col.	Label	Content
A	Species	Identifies the species being modeled by four-letter code.
B	Link#	Contains a unique identification number for each causal link.
C	Life Stage	Identifies the life stage affected by the link.
D	Causal Node Type	Identifies whether the causal node for the link is a controlling factor, habitat element, critical biological activity or process, or life-stage outcome.
E	Causal Node	Identifies the causal node in the link.
F	Effect Node Type	Identifies whether the effect node for the link is a controlling factor, habitat element, critical biological activity or process, or life-stage outcome.
G	Effect Node	Identifies the effect node in the link.
H	Link Reason	States the rationale for including the link in the conceptual ecological model, including citations as appropriate.
I	Link Character Type	Identifies the character of the link based on standard definitions.
J	Link Character Direction	Identifies whether the link is uni- or bi-directional.
K	Link Character Reason	States the rationale for the entries for Link Character Type and Link Character Direction, including citations as appropriate.
L	Link Intensity	Shows the rating of link intensity based on the definitions in table 1-1.
M	Link Spatial Scale	Shows the rating of link spatial scale based on the definitions in table 1-2.
N	Link Temporal Scale	Shows the rating of link temporal scale based on the definitions in table 1-3.
O	Link Average Magnitude	Shows the numerical average rating of link intensity, spatial scale, and temporal scale based on the definitions in table 1-4.
P	Link Magnitude Rank	Shows the overall rating of link magnitude based on the Link Average Magnitude, grouped following the criteria in table 1-4.
Q	Link Magnitude Reason	States the rationale for the ratings for link intensity, spatial scale, and temporal scale, with citations as appropriate.
R	Link Predictability Rank	Shows the rating of link predictability based on the definitions in table 1-5.
S	Link Predictability Reason	States the rationale for the rating of link predictability, with citations as appropriate.
T	Link Understanding Rank	Shows the rating of link understanding based on the definitions in table 1-6.
U	Link Understanding Reason	States the rationale for the rating of link predictability, including comments on alternative interpretations and publications/experts associated with different interpretations when feasible, with citations as appropriate.
V	Management Questions	Briefly notes questions that appear to arise from the preceding entries for the link, focused on critical gaps or uncertainties in knowledge concerning <i>management actions and options</i> , with reasoning, including the estimate of relative importance when possible.
W	Research Questions	Brief notes that appear to arise from the preceding entries for the link, focused on critical gaps or uncertainties in <i>basic scientific knowledge</i> , with reasoning, including the estimate of relative importance when possible.
X	Other Comments	Provides additional notes on investigator concerns, uncertainties, and questions.
Y	Update Status	Provides information on the history of editing the information on this link for updates carried out after completion of an initial version.




Figure 1-1.—Conventions for displaying cause and effect nodes, linkages, link magnitude, link understanding, and link predictability.

## Literature Cited

- Burke, M., K. Jorde, and J.M. Buffington. 2009. Application of a hierarchical framework for assessing environmental impacts of dam operation: changes in streamflow, bed mobility and recruitment of riparian trees in a western North American river. *Journal of Environmental Management* 90:S224–S236.
- DiGennaro, B., D. Reed, C. Swanson, L. Hastings, Z. Hymanson, M. Healey, S. Siegel, S. Cantrell, and B. Herbold. 2012. Using conceptual models and decision-support tools to guide ecosystem restoration planning and adaptive management: an example from the Sacramento–San Joaquin Delta, California. *San Francisco Estuary and Watershed Science* 10(3):1–15. <http://escholarship.org/uc/item/3j95x7vt>
- Fischenich, J.C. 2008. The application of conceptual models to ecosystem restoration. U.S. Army Corps of Engineers, Engineer Research and Development Center (ERDC), Ecosystem Management and Restoration Research Program (EMRRP), Technical Note ERDC/EBA TN-08-1, February 2008. Vicksburg, Mississippi. <http://el.ercd.usace.army.mil/publications.cfm?Topic=technote&Code=emrrp>
- Kondolf, G.M., J.G. Williams, T.C. Horner, and D. Milan. 2008. Assessing physical quality of spawning habitat. Pages 249–274 *in* D.A. Sear and P. DeVries (editors). *Salmonid Spawning Habitat in Rivers: Physical Controls, Biological Responses, and Approaches*. American Fisheries Society Symposium 65. American Fisheries Society, Bethesda, Maryland.
- McDonald, D.B. and H. Caswell. 1993. Matrix methods for avian demography. Pages 139–185 *in* D.M. Power (editor). *Current Ornithology*. Plenum Press: New York, New York.
- Wildhaber, M.L., A.J. DeLonay, D.M. Papoulias, D.L. Galat, R.B. Jacobson, D.G. Simpkins, P.J. Baaten, C.E. Korschgen, and M.J. Mac. 2007. A conceptual life-history model for pallid and shovelnose sturgeon. U.S. Geological Survey, Circular 1315. Reston, Virginia.
- \_\_\_\_\_. 2011. Identifying structural elements needed for development of a predictive life-history model for pallid and shovelnose sturgeons. *Journal of Applied Ichthyology* 27:462–469.

## **ATTACHMENT 2**

Vermilion Flycatcher Habitat Data

<b>Habitat element</b>	<b>Value or range</b>	<b>Location</b>	<b>Reference</b>
Canopy cover	Not quantified		
Patch size	Not quantified		

Note: The data presented in this table reflect those available in the literature at the time this model was developed. These data have not been validated.